

ON THE SPECTRAL ASYMPTOTICS OF OPERATORS ON MANIFOLDS WITH ENDS

SANDRO CORIASCO AND LIDIA MANICCIA

ABSTRACT. We deal with the asymptotic behaviour for $\lambda \rightarrow +\infty$ of the counting function $N_P(\lambda)$ of certain positive selfadjoint operators P with double order (m, μ) , $m, \mu > 0$, $m \neq \mu$, defined on a manifold with ends M . The structure of this class of noncompact manifolds allows to make use of calculi of pseudodifferential operators and Fourier Integral Operators associated with weighted symbols globally defined on \mathbb{R}^n . By means of these tools, we improve known results concerning the remainder terms of the Weyl Formulae for $N_P(\lambda)$ and show how their behaviour depends on the ratio $\frac{m}{\mu}$ and the dimension of M .

1. INTRODUCTION

The aim of this paper is to study the asymptotic behaviour, for $\lambda \rightarrow +\infty$, of the counting function

$$N_P(\lambda) = \sum_{\lambda_j \leq \lambda} 1$$

where $\lambda_1 \leq \lambda_2 \leq \dots$ is the sequence of the eigenvalues, repeated according to their multiplicities, of a positive order, selfadjoint, classical, elliptic SG-pseudodifferential operator P on a manifold with ends. Explicitly, SG-pseudodifferential operators $P = p(x, D) = \text{Op}(p)$ on \mathbb{R}^n can be defined via the usual left-quantization

$$Pu(x) = \frac{1}{(2\pi)^n} \int e^{ix \cdot \xi} p(x, \xi) \hat{u}(\xi) d\xi, \quad u \in \mathcal{S}(\mathbb{R}^n),$$

starting from symbols $p(x, \xi) \in C^\infty(\mathbb{R}^n \times \mathbb{R}^n)$ with the property that, for arbitrary multiindices α, β , there exist constants $C_{\alpha\beta} \geq 0$ such that the estimates

$$(1.1) \quad |D_\xi^\alpha D_x^\beta p(x, \xi)| \leq C_{\alpha\beta} \langle \xi \rangle^{m-|\alpha|} \langle x \rangle^{\mu-|\beta|}$$

hold for fixed $m, \mu \in \mathbb{R}$ and all $(x, \xi) \in \mathbb{R}^n \times \mathbb{R}^n$, where $\langle y \rangle = \sqrt{1 + |y|^2}$, $y \in \mathbb{R}^n$. Symbols of this type belong to the class denoted by $S^{m,\mu}(\mathbb{R}^n)$, and the corresponding operators constitute the class $L^{m,\mu}(\mathbb{R}^n) = \text{Op}(S^{m,\mu}(\mathbb{R}^n))$. In the sequel we will sometimes write $S^{m,\mu}$ and $L^{m,\mu}$, respectively, fixing once and for all the dimension of the (non-compact) base manifold to n .

These classes of operators, introduced on \mathbb{R}^n by H.O. Cordes [6] and C. Parenti [30], see also R. Melrose [26], M.A. Shubin [33], form a graded algebra, i.e., $L^{r,\rho} \circ L^{m,\mu} \subseteq L^{r+m,\rho+\mu}$. The remainder elements are operators with symbols in $S^{-\infty,-\infty}(\mathbb{R}^n) = \bigcap_{(m,\mu) \in \mathbb{R}^2} S^{m,\mu}(\mathbb{R}^n) = \mathcal{S}(\mathbb{R}^{2n})$, that is, those having kernel in $\mathcal{S}(\mathbb{R}^{2n})$, con-

tinuously mapping $\mathcal{S}'(\mathbb{R}^n)$ to $\mathcal{S}(\mathbb{R}^n)$. An operator $P = \text{Op}(p) \in L^{m,\mu}$ and its symbol $p \in S^{m,\mu}$ are called SG-elliptic if there exists $R \geq 0$ such that $p(x, \xi)$ is invertible for $|x| + |\xi| \geq R$ and

$$p(x, \xi)^{-1} = O(\langle \xi \rangle^{-m} \langle x \rangle^{-\mu}).$$

2010 *Mathematics Subject Classification.* Primary: 58J40; Secondary: 35S05, 35S30, 47G30, 58J45.
Key words and phrases. Manifold with ends, Spectral asymptotics, Weyl formula.

In such case we will usually write $P \in EL^{m,\mu}$. Operators in $L^{m,\mu}$ act continuously from $\mathcal{S}(\mathbb{R}^n)$ to itself, and extend as continuous operators from $\mathcal{S}'(\mathbb{R}^n)$ to itself and from $H^{s,\sigma}(\mathbb{R}^n)$ to $H^{s-m,\sigma-\mu}(\mathbb{R}^n)$, where $H^{s,\sigma}(\mathbb{R}^n)$, $s, \sigma \in \mathbb{R}$, denotes the weighted Sobolev space

$$H^{s,\sigma}(\mathbb{R}^n) = \{u \in \mathcal{S}'(\mathbb{R}^n) : \|u\|_{s,\sigma} = \|\text{Op}(\pi_{s,\sigma})u\|_{L^2} < \infty\},$$

$$\pi_{s,\sigma}(x, \xi) = \langle \xi \rangle^s \langle x \rangle^\sigma.$$

Continuous inclusions $H^{s,\sigma}(\mathbb{R}^n) \hookrightarrow H^{r,\rho}(\mathbb{R}^n)$ hold when $s \geq r$ and $\sigma \geq \rho$, compact when both inequalities are strict, and

$$\mathcal{S}(\mathbb{R}^n) = \bigcap_{(s,\sigma) \in \mathbb{R}^2} H^{s,\sigma}(\mathbb{R}^n), \quad \mathcal{S}'(\mathbb{R}^n) = \bigcup_{(s,\sigma) \in \mathbb{R}^2} H^{s,\sigma}(\mathbb{R}^n).$$

An elliptic SG-operator $P \in L^{m,\mu}$ admits a parametrix $E \in L^{-m,-\mu}$ such that

$$PE = I + K_1, \quad EP = I + K_2,$$

for suitable $K_1, K_2 \in L^{-\infty,-\infty} = \text{Op}(\mathcal{S}^{-\infty,-\infty})$, and it turns out to be a Fredholm operator. In 1988, E. Schrohe [31] introduced a class of non-compact manifolds, the so-called SG-manifolds, on which it is possible to transfer from \mathbb{R}^n the whole SG-calculus. In short, these are manifolds which admit a finite atlas whose changes of coordinates behave like symbols of order $(0, 1)$ (see [31] for details and additional technical hypotheses). The manifolds with cylindrical ends are a special case of SG-manifolds, on which also the concept of SG-classical operator makes sense: moreover, the principal symbol of a SG-classical operator P on a manifold with cylindrical ends M , in this case a triple $\sigma(P) = (\sigma_\psi(P), \sigma_e(P), \sigma_{\psi e}(P)) = (p_\psi, p_e, p_{\psi e})$, has an invariant meaning on M , see Y. Egorov and B.-W. Schulze [13], L. Maniccia and P. Panarese [24], R. Melrose [26] and Section 2 below. We indicate the subspaces of classical symbols and operators adding the subscript _{cl} to the notation introduced above.

The literature concerning the study of the eigenvalue asymptotics of elliptic operators is vast, and covers a number of different situations and operator classes, see, e.g., the monograph by V.J. Ivrii [22]. Then, we only mention a few of the many existing papers and books on this deeply investigated subject, which are related to the case we consider here, either by the type of symbols and underlying spaces, or by the techniques which are used: we refer the reader to the corresponding reference lists for more complete informations. On compact manifolds, well known results were proved by L. Hörmander [19] and V. Guillemin [15], see also the book by H. Kumano-go [23]. On the other hand, for operators globally defined on \mathbb{R}^n , see P. Boggiatto, E. Buzano, L. Rodino [2], B. Helffer [16], L. Hörmander [20], A. Mohammed [27], F. Nicola [28], M. A. Shubin [33]. Many other situations have been considered, see the cited book by V.J. Ivrii. On manifolds with ends, T. Christiansen and M. Zworski [5] studied the Laplace-Beltrami operator associated with a scattering metric, while L. Maniccia and P. Panarese [24] applied the heat kernel method to study operators similar to those considered here.

Here we deal with the case of manifolds with ends for $P \in EL_{\text{cl}}^{m,\mu}(M)$, positive and selfadjoint, such that $m, \mu > 0$, $m \neq \mu$, focusing on the (invariant) meaning of the constants appearing in the corresponding Weyl formulae and on achieving a better estimate of the remainder term. Note that the situation we consider here is different from that of the Laplace-Beltrami operator investigated in [5], where continuous spectrum is present as well: in fact, in view of Theorem 3.2, $\text{spec}(P)$ consists only of a sequence of real isolated eigenvalues $\{\lambda_j\}$ with finite multiplicity.

As recalled above, a first result concerning the asymptotic behaviour of $N_P(\lambda)$ for operators including those considered in this paper was proved in [24], giving,

for $\lambda \rightarrow +\infty$,

$$N_P(\lambda) = \begin{cases} C_1 \lambda^{\frac{n}{m}} + o(\lambda^{\frac{n}{m}}) & \text{for } m < \mu \\ C_0^1 \lambda^{\frac{n}{m}} \log \lambda + o(\lambda^{\frac{n}{m}} \log \lambda) & \text{for } m = \mu \\ C_2 \lambda^{\frac{n}{\mu}} + o(\lambda^{\frac{n}{\mu}}) & \text{for } m > \mu. \end{cases}$$

Note that the constants C_1, C_2, C_0^1 above depend only on the principal symbol of P , which implies that they have an invariant meaning on the manifold M , see Sections 2 and 3 below. On the other hand, in view of the technique used there, the remainder terms appeared in the form $o(\lambda^{\frac{n}{\min\{m,\mu\}}})$ and $o(\lambda^{\frac{n}{m}} \log \lambda)$ for $m \neq \mu$ and $m = \mu$, respectively. An improvement in this direction for operators on \mathbb{R}^n had been achieved by F. Nicola [28], who, in the case $m = \mu$ proved that

$$N_P(\lambda) = C_0^1 \lambda^{\frac{n}{m}} \log \lambda + O(\lambda^{\frac{n}{m}}), \lambda \rightarrow +\infty,$$

while, for $m \neq \mu$, showed that the remainder term has the form $O(\lambda^{\frac{n}{\min\{m,\mu\}} - \varepsilon})$ for a suitable $\varepsilon > 0$. A further improvement of these results in the case $m = \mu$ has recently appeared in U. Battisti and S. Coriasco [1], where it has been shown that, for a suitable $\varepsilon > 0$,

$$N_P(\lambda) = C_0^1 \lambda^{\frac{n}{m}} \log \lambda + C_0^2 \lambda^{\frac{n}{m}} + O(\lambda^{\frac{n}{m} - \varepsilon}), \lambda \rightarrow +\infty.$$

Even the constant C_0^2 has an invariant meaning on M , and both C_0^1 and C_0^2 are explicitly computed in terms of trace operators defined on $L_{\text{cl}}^{m,m}(M)$.

In this paper the remainder estimates in the case $m \neq \mu$ are further improved. More precisely, we first consider the power $Q = P^{\frac{1}{\max\{m,\mu\}}}$ of P (see L. Maniccia, E. Schrohe, J. Seiler [25] for the properties of powers of SG-classical operators). Then, by studying the asymptotic behaviour in λ of the trace of the operator $\widehat{\psi}_\lambda(-Q)$, $\psi_\lambda(t) = \psi(t)e^{-it\lambda}$, $\psi \in C_0^\infty(\mathbb{R})$, defined via a Spectral Theorem and approximated in terms of Fourier Integral Operators, we prove the following

Theorem 1.1. *Let M be a manifold with ends of dimension n and let $P \in EL_{\text{cl}}^{m,\mu}(M)$ be a positive selfadjoint operator such that $m, \mu > 0$, $m \neq \mu$, with domain $H^{m,\mu}(M) \hookrightarrow L^2(M)$. Then, the following Weyl formulae hold for $\lambda \rightarrow +\infty$:*

$$(1.2) \quad N_P(\lambda) = \begin{cases} C_1 \lambda^{\frac{n}{m}} + O(\lambda^{\frac{n}{\mu}}) + O(\lambda^{\frac{n}{m} - \frac{1}{\mu}}) = C_1 \lambda^{\frac{n}{m}} + O(\lambda^{\frac{n}{m} - \varepsilon_1}) & \text{for } m < \mu \\ C_2 \lambda^{\frac{n}{\mu}} + O(\lambda^{\frac{n}{m}}) + O(\lambda^{\frac{n}{\mu} - \frac{1}{m}}) = C_2 \lambda^{\frac{n}{\mu}} + O(\lambda^{\frac{n}{\mu} - \varepsilon_2}) & \text{for } m > \mu. \end{cases}$$

where $\varepsilon_1 = \min \left\{ \frac{1}{\mu}, n \left(\frac{1}{m} - \frac{1}{\mu} \right) \right\}$ and $\varepsilon_2 = \min \left\{ \frac{1}{m}, n \left(\frac{1}{\mu} - \frac{1}{m} \right) \right\}$.

The order of the remainder is then determined by the ratio of m and μ and the dimension of M , since

$$(1.3) \quad \begin{aligned} \frac{n}{m} - \frac{1}{\mu} &\leq \frac{n}{\mu} \text{ for } m < \mu \Leftrightarrow 1 < \frac{\mu}{m} \leq 1 + \frac{1}{n}, \\ \frac{n}{\mu} - \frac{1}{m} &\leq \frac{n}{m} \text{ for } m > \mu \Leftrightarrow 1 < \frac{m}{\mu} \leq 1 + \frac{1}{n}. \end{aligned}$$

In particular, when $\frac{\max\{m,\mu\}}{\min\{m,\mu\}} \geq 2$, the remainder is always $O(\lambda^{\frac{n}{\max\{m,\mu\}}})$.

Examples include operators of Schrödinger type on M , that is $P = -\Delta_g + V$, Δ_g the Laplace-Beltrami operator in M associated with a suitable metric g , V a smooth potential that, in the local coordinates $x \in U_N \subseteq \mathbb{R}^n$ on the cylindrical end grows as $\langle x \rangle^\mu$, with an appropriate $\mu > 0$ related to g . Such examples will be discussed in detail, together with the sharpness of the results in Theorem 1.1, in the forthcoming paper [4], see also [3].

The key point in the proof of Theorem 1.1 is the study of the asymptotic behaviour for $\lambda \rightarrow +\infty$ of integrals of the form

$$I(\lambda) = \int e^{i(-t\lambda + \varphi(t;x,\xi) - x\xi)} \psi(t) a(t; x, \xi) dt d\xi dx$$

where a and φ satisfy certain growth conditions in x and ξ (see Section 3 for more details). The integrals $I(\lambda)$ represent in fact the local expressions of the trace of $\widehat{\psi}_\lambda(-Q)$, obtained through the so-called “geometric optic method”, specialised to the SG situation, see e.g. S. Coriasco [7, 8], S. Coriasco and L. Rodino [11]. To treat the integrals $I(\lambda)$ we proceed similarly to A. Grigis and J. Sjöstrand [14], B. Helffer and D. Robert [17], see also H. Tamura [34].

The paper is organised as follows. Section 2 is devoted to recall the definition of SG-classical operators on a manifold with ends M . In Section 3 we show that the asymptotic behaviour of $N_P(\lambda)$, $\lambda \rightarrow +\infty$, for a positive self-adjoint operator $P \in L_{\text{cl}}^{m,\mu}(M)$, $m, \mu > 0$, is related to the asymptotic behaviour of oscillatory integrals of the form $I(\lambda)$. In Section 4 we conclude the proof of Theorem 1.1, investigating the behaviour of $I(\lambda)$ for $\lambda \rightarrow +\infty$. Finally, some technical details are collected in the Appendix.

ACKNOWLEDGEMENTS

The authors wish to thank U. Battisti, L. Rodino and E. Schrohe for useful discussions and hints. The first author gratefully acknowledges the support by the Institut für Analysis, Fakultät für Mathematik und Physik, Gottfried Wilhelm Leibniz Universität Hannover, during his stay as Visiting Scientist in the Academic Year 2011/2012, where this paper was partly developed and completed.

2. SG-CLASSICAL OPERATORS ON MANIFOLDS WITH ENDS

From now on, we will be concerned with the subclass of SG-operators given by those elements $P \in L^{m,\mu}(\mathbb{R}^n)$, $(m, \mu) \in \mathbb{R}^2$, which are SG-classical, that is, $P = \text{Op}(p)$ with $p \in S_{\text{cl}}^{m,\mu}(\mathbb{R}^n) \subset S^{m,\mu}(\mathbb{R}^n)$. We begin recalling the basic definitions and results (see, e.g., [13, 25] for additional details and proofs).

Definition 2.1. i) A symbol $p(x, \xi)$ belongs to the class $S_{\text{cl}(\xi)}^{m,\mu}(\mathbb{R}^n)$ if there exist $p_{m-i,\cdot}(x, \xi) \in \widetilde{\mathcal{H}}_\xi^{m-i}(\mathbb{R}^n)$, $i = 0, 1, \dots$, positively homogeneous functions of order $m - i$ with respect to the variable ξ , smooth with respect to the variable x , such that, for a 0-excision function ω ,

$$p(x, \xi) - \sum_{i=0}^{N-1} \omega(\xi) p_{m-i,\cdot}(x, \xi) \in S^{m-N,\mu}(\mathbb{R}^n), \quad N = 1, 2, \dots;$$

ii) A symbol $p(x, \xi)$ belongs to the class $S_{\text{cl}(x)}^{m,\mu}(\mathbb{R}^n)$ if there exist $p_{\cdot,\mu-k}(x, \xi) \in \widetilde{\mathcal{H}}_x^{\mu-k}(\mathbb{R}^n)$, $k = 0, \dots$, positively homogeneous functions of order $\mu - k$ with respect to the variable x , smooth with respect to the variable ξ , such that, for a 0-excision function ω ,

$$p(x, \xi) - \sum_{k=0}^{N-1} \omega(x) p_{\cdot,\mu-k}(x, \xi) \in S^{m,\mu-N}(\mathbb{R}^n), \quad N = 1, 2, \dots$$

Definition 2.2. A symbol $p(x, \xi)$ is SG-classical, and we write $p \in S_{\text{cl}(x,\xi)}^{m,\mu}(\mathbb{R}^n) = S_{\text{cl}}^{m,\mu}(\mathbb{R}^n) = S_{\text{cl}}^{m,\mu}$, if

- i) there exist $p_{m-j,\cdot}(x, \xi) \in \widetilde{\mathcal{H}}_\xi^{m-j}(\mathbb{R}^n)$ such that, for a 0-excision function ω , $\omega(\xi) p_{m-j,\cdot}(x, \xi) \in S_{\text{cl}(x)}^{m-j,\mu}(\mathbb{R}^n)$ and

$$p(x, \xi) - \sum_{j=0}^{N-1} \omega(\xi) p_{m-j,\cdot}(x, \xi) \in S^{m-N,\mu}(\mathbb{R}^n), \quad N = 1, 2, \dots;$$

- ii) there exist $p_{\cdot,\mu-k}(x, \xi) \in \widetilde{\mathcal{H}}_x^{\mu-k}(\mathbb{R}^n)$ such that, for a 0-excision function ω , $\omega(x) p_{\cdot,\mu-k}(x, \xi) \in S_{\text{cl}(\xi)}^{m,\mu-k}(\mathbb{R}^n)$ and

$$p(x, \xi) - \sum_{k=0}^{N-1} \omega(x) p_{\cdot,\mu-k}(x, \xi) \in S^{m,\mu-N}(\mathbb{R}^n), \quad N = 1, 2, \dots$$

We set $L_{\text{cl}(x,\xi)}^{m,\mu}(\mathbb{R}^n) = L_{\text{cl}}^{m,\mu} = \text{Op}(S_{\text{cl}}^{m,\mu})$.

Remark 2.3. The definition could be extended in a natural way from operators acting between scalars to operators acting between (distributional sections of) vector bundles: one should then use matrix-valued symbols whose entries satisfy the estimates (1.1).

Note that the definition of SG-classical symbol implies a condition of compatibility for the terms of the expansions with respect to x and ξ . In fact, defining σ_ψ^{m-j} and $\sigma_e^{\mu-i}$ on $S_{\text{cl}(\xi)}^{m,\mu}$ and $S_{\text{cl}(x)}^{m,\mu}$, respectively, as

$$\begin{aligned} \sigma_\psi^{m-j}(p)(x, \xi) &= p_{m-j,\cdot}(x, \xi), \quad j = 0, 1, \dots, \\ \sigma_e^{\mu-i}(p)(x, \xi) &= p_{\cdot,\mu-i}(x, \xi), \quad i = 0, 1, \dots, \end{aligned}$$

it possible to prove that

$$p_{m-j,\mu-i} = \sigma_{\psi e}^{m-j,\mu-i}(p) = \sigma_\psi^{m-j}(\sigma_e^{\mu-i}(p)) = \sigma_e^{\mu-i}(\sigma_\psi^{m-j}(p)),$$

$$j = 0, 1, \dots, i = 0, 1, \dots$$

Moreover, the composition of two SG-classical operators is still classical. For $P = \text{Op}(p) \in L_{\text{cl}}^{m,\mu}$ the triple $\sigma(P) = (\sigma_\psi(P), \sigma_e(P), \sigma_{\psi e}(P)) = (p_{m,\cdot}, p_{\cdot,\mu}, p_{m,\mu}) = (p_\psi, p_e, p_{\psi e})$ is called the *principal symbol* of P . The three components are also called the ψ -, e - and ψe -principal symbol, respectively. This definition keeps the usual multiplicative behaviour, that is, for any $R \in L_{\text{cl}}^{r,\rho}$, $S \in L_{\text{cl}}^{s,\sigma}$, $(r, \rho), (s, \sigma) \in \mathbb{R}^2$, $\sigma(RS) = \sigma(S) \sigma(R)$, with componentwise product in the right-hand side. We also set

$$\begin{aligned} \text{Sym}_p(P)(x, \xi) &= \text{Sym}_p(p)(x, \xi) = \\ &= p_{\mathbf{m}}(x, \xi) = \omega(\xi) p_\psi(x, \xi) + \omega(x) (p_e(x, \xi) - \omega(\xi) p_{\psi e}(x, \xi)), \end{aligned}$$

for a fixed 0-excision function ω . Theorem 2.4 below allows to express the ellipticity of SG-classical operators in terms of their principal symbol:

Theorem 2.4. An operator $P \in L_{\text{cl}}^{m,\mu}$ is elliptic if and only if each element of the triple $\sigma(P)$ is non-vanishing on its domain of definition.

As a consequence, denoting by $\{\lambda_j\}$ the sequence of eigenvalues of P , ordered such that $j \leq k \Rightarrow \lambda_j \leq \lambda_k$, with each eigenvalue repeated accordingly to its multiplicity, the counting function $N_P(\lambda) = \sum_{\lambda_j \leq \lambda} 1$ is well-defined for a SG-classical elliptic self-

adjoint operator P , see, e.g., [1, 3, 4, 28]. We now introduce the class of noncompact manifolds with which we will deal:

Definition 2.5. A manifold with a cylindrical end is a triple $(M, X, [f])$, where $M = \mathcal{M} \amalg_{\mathcal{C}} \mathcal{C}$ is a n -dimensional smooth manifold and

- i) \mathcal{M} is a smooth manifold, given by $\mathcal{M} = (M_0 \setminus D) \cup C$ with a n -dimensional smooth compact manifold without boundary M_0 , D a closed disc of M_0 and $C \subset D$ a collar neighbourhood of ∂D in M_0 ;
- ii) \mathcal{C} is a smooth manifold with boundary $\partial\mathcal{C} = X$, with X diffeomorphic to ∂D ;
- iii) $f : [\delta_f, \infty) \times \mathbb{S}^{n-1} \rightarrow \mathcal{C}$, $\delta_f > 0$, is a diffeomorphism, $f(\{\delta_f\} \times \mathbb{S}^{n-1}) = X$ and $f([\delta_f, \delta_f + \varepsilon_f] \times \mathbb{S}^{n-1})$, $\varepsilon_f > 0$, is diffeomorphic to C ;
- iv) the symbol Π_C means that we are gluing \mathcal{M} and \mathcal{C} , through the identification of C and $f([\delta_f, \delta_f + \varepsilon_f] \times \mathbb{S}^{n-1})$;
- v) the symbol $[f]$ represents an equivalence class in the set of functions

$g : [\delta_g, \infty) \times \mathbb{S}^{n-1} \rightarrow \mathcal{C}$: g is a diffeomorphism,

$$g(\{\delta_g\} \times \mathbb{S}^{n-1}) = X \text{ and}$$

$$g([\delta_g, \delta_g + \varepsilon_g] \times \mathbb{S}^{n-1}), \varepsilon_g > 0, \text{ is diffeomorphic to } C$$

where $f \sim g$ if and only if there exists a diffeomorphism $\Theta \in \text{Diff}(\mathbb{S}^{n-1})$ such that

$$(2.1) \quad (g^{-1} \circ f)(\rho, \gamma) = (\rho, \Theta(\gamma))$$

for all $\rho \geq \max\{\delta_f, \delta_g\}$ and $\gamma \in \mathbb{S}^{n-1}$.

We use the following notation:

- $U_{\delta_f} = \{x \in \mathbb{R}^n : |x| > \delta_f\}$;
- $\mathcal{C}_\tau = f([\tau, \infty) \times \mathbb{S}^{n-1})$, where $\tau \geq \delta_f$. The equivalence condition (2.1) implies that \mathcal{C}_τ is well defined;
- $\pi : \mathbb{R}^n \setminus \{0\} \rightarrow (0, \infty) \times \mathbb{S}^{n-1} : x \mapsto \pi(x) = (|x|, \frac{x}{|x|})$;
- $f_\pi = f \circ \pi : \overline{U_{\delta_f}} \rightarrow \mathcal{C}$ is a parametrisation of the end. Let us notice that, setting $F = g_\pi^{-1} \circ f_\pi$, the equivalence condition (2.1) implies

$$(2.2) \quad F(x) = |x| \Theta\left(\frac{x}{|x|}\right).$$

We also denote the restriction of f_π mapping U_{δ_f} onto $\mathcal{C} = \mathcal{C} \setminus X$ by \dot{f}_π .

The couple $(\mathcal{C}, \dot{f}_\pi^{-1})$ is called the exit chart. If $\mathcal{A} = \{(\Omega_i, \psi_i)\}_{i=1}^N$ is such that the subset $\{(\Omega_i, \psi_i)\}_{i=1}^{N-1}$ is a finite atlas for \mathcal{M} and $(\Omega_N, \psi_N) = (\mathcal{C}, \dot{f}_\pi^{-1})$, then M , with the atlas \mathcal{A} , is a SG-manifold (see [33]): an atlas \mathcal{A} of such kind is called *admissible*. From now on, we restrict the choice of atlases on M to the class of admissible ones. We introduce the following spaces, endowed with their natural topologies:

$$\mathcal{S}(U_\delta) = \left\{ u \in C^\infty(U_\delta) : \forall \alpha, \beta \in \mathbb{N}^n \forall \delta' > \delta \sup_{x \in U_{\delta'}} |x^\alpha \partial^\beta u(x)| < \infty \right\},$$

$$\mathcal{S}_0(U_\delta) = \bigcap_{\delta' \searrow \delta} \{u \in \mathcal{S}(U_{\delta'}) : \text{supp } u \subseteq \overline{U_{\delta'}}\},$$

$$\mathcal{S}(M) = \{u \in C^\infty(M) : u \circ \dot{f}_\pi \in \mathcal{S}(U_{\delta_f}) \text{ for any exit map } \dot{f}_\pi\},$$

$\mathcal{S}'(M)$ denotes the dual space of $\mathcal{S}(M)$.

Definition 2.6. The set $S^{m,\mu}(U_{\delta_f})$ consists of all the symbols $a \in C^\infty(U_{\delta_f})$ which fulfill (1.1) for $(x, \xi) \in U_{\delta_f} \times \mathbb{R}^n$ only. Moreover, the symbol a belongs to the subset $SG_{\text{cl}}^{m,\mu}(U_{\delta_f})$ if it admits expansions in asymptotic sums of homogeneous symbols with respect to x and ξ as in Definitions 2.1 and 2.2, where the remainders are now given by SG-symbols of the required order on U_{δ_f} .

Note that, since U_{δ_f} is conical, the definition of homogeneous and classical symbol on U_{δ_f} makes sense. Moreover, the elements of the asymptotic expansions of the

classical symbols can be extended by homogeneity to smooth functions on $\mathbb{R}^n \setminus \{0\}$, which will be denoted by the same symbols. It is a fact that, given an admissible atlas $\{(\Omega_i, \psi_i)\}_{i=1}^N$ on M , there exists a partition of unity $\{\theta_i\}$ and a set of smooth functions $\{\chi_i\}$ which are compatible with the SG-structure of M , that is:

- $\text{supp } \theta_i \subset \Omega_i, \text{supp } \chi_i \subset \Omega_i, \chi_i \theta_i = \theta_i, i = 1, \dots, N;$
- $|\partial^\alpha(\theta_N \circ f_\pi)(x)| \leq C_\alpha \langle x \rangle^{-|\alpha|}$ and $|\partial^\alpha(\chi_N \circ f_\pi)(x)| \leq C_\alpha \langle x \rangle^{-|\alpha|}$ for all $x \in U_{\delta_f}$.

Moreover, θ_N and χ_N can be chosen so that $\theta_N \circ f_\pi$ and $\chi_N \circ f_\pi$ are homogeneous of degree 0 on U_δ . We denote by u^* the composition of $u: \psi_i(\Omega_i) \subset \mathbb{R}^n \rightarrow \mathbb{C}$ with the coordinate patches ψ_i , and by v_* the composition of $v: \Omega_i \subset M \rightarrow \mathbb{C}$ with $\psi_i^{-1}, i = 1, \dots, N$. It is now possible to give the definition of SG-pseudodifferential operator on M :

Definition 2.7. Let M be a manifold with a cylindrical end. A linear operator $P: \mathcal{S}(M) \rightarrow \mathcal{S}'(M)$ is a SG-pseudodifferential operator of order (m, μ) on M , and we write $P \in L^{m, \mu}(M)$, if, for any admissible atlas $\{(\Omega_i, \psi_i)\}_{i=1}^N$ on M with exit chart (Ω_N, ψ_N) :

- 1) for all $i = 1, \dots, N-1$ and any $\theta_i, \chi_i \in C_c^\infty(\Omega_i)$, there exist symbols $p^i(x, \xi) \in S^m(\psi_i(\Omega_i))$ such that

$$(\chi_i P \theta_i u^*)_*(x) = \iint e^{i(x-y) \cdot \xi} p^i(x, \xi) u(y) dy dx, \quad u \in C^\infty(\psi_i(\Omega_i));$$

- 2) for any θ_N, χ_N of the type described above, there exists a symbol $p^N(x, \xi) \in SG^{m, \mu}(U_{\delta_f})$ such that

$$(\chi_N P \theta_N u^*)_*(x) = \iint e^{i(x-y) \cdot \xi} p^N(x, \xi) u(y) dy dx, \quad u \in \mathcal{S}_0(U_{\delta_f});$$

- 3) K_P , the Schwartz kernel of P , is such that

$$K_P \in C^\infty((M \times M) \setminus \Delta) \bigcap \mathcal{S}((\mathcal{C} \times \mathcal{C}) \setminus W)$$

where Δ is the diagonal of $M \times M$ and $W = (f_\pi \times f_\pi)(V)$ with any conical neighbourhood V of the diagonal of $U_{\delta_f} \times U_{\delta_f}$.

The most important local symbol of P is p^N . Our definition of SG-classical operator on M differs slightly from the one in [24]:

Definition 2.8. Let $P \in L^{m, \mu}(M)$. P is a SG-classical operator on M , and we write $P \in L_{\text{cl}}^{m, \mu}(M)$, if $p^N(x, \xi) \in S_{\text{cl}}^{m, \mu}(U_{\delta_f})$ and the operator P , restricted to the manifold \mathcal{M} , is classical in the usual sense.

The usual homogeneous principal symbol p_ψ of a SG-classical operator $P \in L_{\text{cl}}^{m, \mu}(M)$ is of course well-defined as a smooth function on T^*M . In order to give an invariant definition of the principal symbols homogeneous in x of an operator $P \in L_{\text{cl}}^{m, \mu}(M)$, the subbundle $T_X^*M = \{(x, \xi) \in T^*M: x \in X, \xi \in T_x^*M\}$ was introduced. The notions of ellipticity can be extended to operators on M as well:

Definition 2.9. Let $P \in L_{\text{cl}}^{m, \mu}(M)$ and let us fix an exit map f_π . We can define local objects $p_{m-j, \mu-i}, p_{\cdot, \mu-i}$ as

$$\begin{aligned} p_{m-j, \mu-i}(\theta, \xi) &= p_{m-j, \mu-i}^N(\theta, \xi), \quad \theta \in \mathbb{S}^{n-1}, \xi \in \mathbb{R}^n \setminus \{0\}, \\ p_{\cdot, \mu-i}(\theta, \xi) &= p_{\cdot, \mu-i}^N(\theta, \xi), \quad \theta \in \mathbb{S}^{n-1}, \xi \in \mathbb{R}^n. \end{aligned}$$

Definition 2.10. An operator $P \in L_{\text{cl}}^{m, \mu}(M)$ is elliptic, and we write $P \in EL_{\text{cl}}^{m, \mu}(M)$, if the principal part of $p^N \in S^{m, \mu}(U_{\delta_f})$ satisfies the SG-ellipticity conditions on $U_{\delta_f} \times \mathbb{R}^n$ and the operator P , restricted to the manifold \mathcal{M} , is elliptic in the usual sense.

Proposition 2.11. *The properties $P \in L^{m,\mu}_{\text{cl}}(M)$ and $P \in L^{m,\mu}_{\text{cl}}(M)$, as well as the notion of SG-ellipticity, do not depend on the (admissible) atlas on M . Moreover, the local functions p_e and p_{ψ_e} give rise to invariantly defined elements of $C^\infty(T^*_X M)$ and $C^\infty(T^*_X M \setminus 0)$, respectively.*

Then, with any $P \in L^{m,\mu}_{\text{cl}}(M)$, it is associated an invariantly defined principal symbol in three components $\sigma(P) = (p_\psi, p_e, p_{\psi_e})$. Finally, through local symbols given by $\pi^j(x, \xi) = \langle \xi \rangle^s$, $j = 1, \dots, N-1$, and $\pi^N(x, \xi) = \langle \xi \rangle^s \langle x \rangle^\sigma$, $s, \sigma \in \mathbb{R}$, we get a SG-elliptic operator $\Pi_{s,\sigma} \in L^{s,\sigma}_{\text{cl}}(M)$ and introduce the (invariantly defined) weighted Sobolev spaces $H^{s,\sigma}(M)$ as

$$H^{s,\sigma}(M) = \{u \in \mathcal{S}'(M) : \Pi_{s,\sigma} u \in L^2(M)\}.$$

The properties of the spaces $H^{s,\sigma}(\mathbb{R}^n)$ extend to $H^{s,\sigma}(M)$ without any change, as well as the continuity of the linear mappings $P: H^{s,\sigma}(M) \rightarrow H^{s-m,\sigma-\mu}(M)$ induced by $P \in L^{m,\mu}_{\text{cl}}(M)$, mentioned in Section 1.

3. SPECTRAL ASYMPTOTICS FOR SG-CLASSICAL ELLIPTIC SELF-ADJOINT OPERATORS ON MANIFOLDS WITH ENDS

In this section we illustrate the procedure to prove Theorem 1.1, similarly to [14], [16], [34]. The result will follow from the Trace formula (3.6), (3.7), the asymptotic behaviour (3.8) and the Tauberian Theorem 3.6. The remaining technical points, in particular the proof of the asymptotic behaviour of the integrals appearing in (3.7), are described in Section 4 and in the Appendix.

Let the operator $P \in EL^{m,\mu}_{\text{cl}}(M)$ be considered as an unbounded operator $P: \mathcal{S}(M) \subset H^{0,0}(M) = L^2(M) \rightarrow L^2(M)$. The following Proposition can be proved by reducing to the local situation and using continuity and ellipticity of P , its parametrix and the density of $\mathcal{S}(M)$ in the $H^{s,\sigma}(M)$ spaces,

Proposition 3.1. *Every $P \in EL^{m,\mu}_{\text{cl}}(M)$, considered as an unbounded operator $P: \mathcal{S}(M) \subset L^2(M) \rightarrow L^2(M)$, admits a unique closed extension, still denoted by P , whose domain is $\mathcal{D}(P) = H^{m,\mu}(M)$.*

From now on, when we write $P \in EL^{m,\mu}_{\text{cl}}(M)$ we always mean its unique closed extension, defined in Proposition 3.1. As standard, we denote by $\rho(P)$ the resolvent set of P , i.e., the set of all $\lambda \in \mathbb{C}$ such that $\lambda I - P$ maps $H^{m,\mu}(M)$ bijectively onto $L^2(M)$. The spectrum of P is then $\text{spec}(P) = \mathbb{C} \setminus \rho(P)$. The next Theorem was proved in [24].

Theorem 3.2. (Spectral Theorem) *Let $P \in EL^{m,\mu}_{\text{cl}}(M)$ be regarded as a closed unbounded operator on $L^2(M)$ with dense domain $H^{m,\mu}(M)$. Assume also that $m, \mu > 0$ and $P^* = P$. Then:*

- i) $(\lambda I - P)^{-1}$ is a compact operator on $L^2(M)$ for every $\lambda \in \rho(P)$. More precisely, $(\lambda I - P)^{-1}$ is an extension by continuity from $\mathcal{S}(M)$ or a restriction from $\mathcal{S}'(M)$ of an operator in $EL^{-m,-\mu}_{\text{cl}}(M)$.
- ii) $\text{spec}(P)$ consists of a sequence of real isolated eigenvalues $\{\lambda_j\}$ with finite multiplicity, clustering at infinity; the orthonormal system of eigenfunctions $\{e_j\}_{j \geq 1}$ is complete in $L^2(M) = H^{0,0}(M)$. Moreover, $e_j \in \mathcal{S}(M)$ for all j .

Given a positive selfadjoint operator $P \in EL^{m,\mu}_{\text{cl}}(M)$, $m, \mu > 0$, $\mu \neq m$, we can assume, without loss of generality¹, $1 \leq \lambda_1 \leq \lambda_2 \dots$. Define the counting function $N_P(\lambda)$, $\lambda \in \mathbb{R}$, as

$$(3.1) \quad N_P(\lambda) = \sum_{\lambda_j \leq \lambda} 1 = \#(\text{spec}(P) \cap (-\infty, \lambda]).$$

¹Considering, if necessary, $P + c$ in place of P , with $c \in \mathbb{R}$ a suitably large constant.

Clearly, N_P is non-decreasing, continuous from the right and supported in $[0, +\infty)$. If we set $Q = P^\dagger$, $l = \max\{m, \mu\}$ (see [25] for the definition of the powers of P), Q turns out to be a SG-classical elliptic selfadjoint operator with $\sigma(Q) = (p_{\psi'}^\dagger, p_e^\dagger, p_{\psi_e}^\dagger)$.

We denote by $\{\eta_j\}$ the sequence of eigenvalues of Q , which satisfy $\eta_j = \lambda_j^{\frac{1}{l}}$: we can then, as above, consider $N_Q(\eta)$. It is a fact that $N_Q(\eta) = O(\eta^{\frac{n}{l}})$, see [24].

From now on we focus on the case $\mu > m > 0$: the case $m > \mu > 0$ can be treated in a completely similar way, exchanging the role of x and ξ . So we can start from a closed positive selfadjoint operator $Q \in EL_{\text{cl}}^{m,1}(M)$ with domain $\mathcal{D}(Q) = H^{m,1}(M)$, $m \in (0, 1)$. For $u \in H^{m,1}(M)$, $t \in \mathbb{R}$, we set

$$(3.2) \quad U(t)u = \sum_{j=1}^{\infty} e^{it\eta_j} (u, e_j)_{L^2(M)} e_j,$$

and the series converges in the $L^2(M)$ norm (cfr., e.g., [14]). Clearly, for all $t \in \mathbb{R}$, $U(t)$ is a unitary operator such that

$$U(0) = I, \quad U(t+s) = U(t)U(s), \quad t, s \in \mathbb{R}.$$

Moreover, if $u \in H^{km,k}(M)$ for some $k \in \mathbb{N}$, $U(t)u \in C^k(\mathbb{R}, H^{0,0}(M)) \cap \dots \cap C^0(\mathbb{R}, H^{km,k}(M))$ and, for $u \in H^{m,1}(M)$, we have $D_t U(t)u - QU(t)u = 0$, $U(0)u = u$, which implies that $v(t, x) = U(t)u(x)$ is a solution of the Cauchy problem

$$(3.3) \quad (D_t - Q)v = 0, \quad v|_{t=0} = u.$$

Let us fix $\psi \in \mathcal{S}(\mathbb{R})$. We can then define the operator $\widehat{\psi}(-Q)$ either by using the formula

$$\widehat{\psi}(-Q)u = \sum_{j=1}^{\infty} \widehat{\psi}(-\eta_j) (u, e_j)_{L^2(M)} e_j$$

or by means of the vector-valued integral $\left(\int \psi(t) U(t) dt \right) u = \int \psi(t) U(t) u dt$, $u \in$

$H^{0,0}(M)$. Indeed, there exists $N_0 \in \mathbb{N}$ such that $\sum_{j=1}^{\infty} \eta_j^{-N_0} < \infty$, so the definition

makes sense and gives an operator in $\mathcal{L}(L^2(M))$ with norm bounded by $\|\psi\|_{L^1(\mathbb{R})}$. The following Lemma, whose proof can be found in the Appendix, is an analog on M of Proposition 1.10.11 in [16]:

Lemma 3.3. $\widehat{\psi}(-Q)$ is an operator with kernel $K_\psi(x, y) = \sum_j \widehat{\psi}(-\eta_j) e_j(x) \overline{e_j(y)} \in \mathcal{S}(M \times M)$.

Clearly, we then have

$$(3.4) \quad \int_M K_\psi(x, x) dx = \sum_j \widehat{\psi}(-\eta_j).$$

By the analysis in [7, 8], [10], [11] (see also [9]), the above Cauchy Problem (3.3) can be solved modulo $\mathcal{S}(M)$ by means of a smooth family of operators $V(t)$, defined for $t \in (-T, T)$, $T > 0$ suitably small, in the sense that $(D_t - Q) \circ V$ is a family of smoothing operators and $V(0)$ is the identity on $\mathcal{S}'(M)$. More explicitly, the following theorem holds²:

²See the Appendix for some details concerning the extension to the manifold M of the results on \mathbb{R}^n proved in [7, 8], [10], [11].

Theorem 3.4. Define $V(t)u = \sum_{k=1}^N \chi_k A_k(t)(\theta_k u)$, where θ_k and χ_k are as in Definition 2.7, with $\chi_k \theta_k = \theta_k$, $k = 1, \dots, N$, while the $A_k(t)$ are SG FIOs which, in the local coordinate open set $U_k = \psi_k(\Omega_k)$ and with $v \in \mathcal{S}(\mathbb{R}^n)$, are given by

$$(A_k(t)v)(x) = \int e^{i\varphi_k(t;x,\xi)} a_k(t;x,\xi) \hat{v}(\xi) d\xi.$$

Each $A_k(t)$ solves a local Cauchy Problem³ $(D_t - Q_k) \circ A_k \in C^\infty((-T, T), L^{-\infty, -\infty}(\mathbb{R}^n))$, $A_k(0) = I$, with $Q_k = \text{op}(q_k)$ and $\{q_k\} \subset SG_{\text{cl}}^{m,1}(\mathbb{R}^n)$ local (complete) symbol of Q associated with $\{\theta_k\}$, $\{\chi_k\}$, with phase and amplitude functions such that

$$(3.5) \quad \begin{aligned} \partial_t \varphi_k(t; x, \xi) - q_k(x, d_x \varphi_k(t; x, \xi)) &= 0, \varphi_k(0; x, \xi) = x\xi, \\ a_k &\in C^\infty((-T, T), SG_{\text{cl}}^{0,0}(\mathbb{R}^n)), a_k(0; x, \xi) = 1. \end{aligned}$$

Then, $V(t)$ satisfies

$$(D_t - Q) \circ V \in C^\infty((-T, T), L^{-\infty, -\infty}(M)), \quad V(0) = I,$$

and $U - V \in C^\infty((-T, T), L^{-\infty, -\infty}(M))$.

Remark 3.5. Notation like $b \in C^\infty((-T, T), S^{r,\rho}(\mathbb{R}^n))$, $B \in C^\infty((-T, T), L^{r,\rho}(M))$, and similar, in Theorem 3.4 and in the sequel, also mean that the seminorms of the involved elements in the corresponding spaces (induced, in the mentioned cases, by (1.1)), are uniformly bounded with respect to $t \in (-T, T)$.

If we write $\psi_\lambda(t) = \psi(t)e^{-it\lambda}$ in place of $\psi(t)$, for a chosen $\psi \in C_0^\infty((-T, T))$, the trace formula (3.4) becomes

$$(3.6) \quad \int_M K_{\psi_\lambda}(x, x) dx = \sum \widehat{\psi}(\lambda - \eta_j).$$

Let us denote the kernel of $U - V$ by $r(t; x, y) \in C^\infty((-T, T), \mathcal{S}(M \times M))$. Then, the distribution kernel of $\int e^{-it\lambda} \psi(t) U(t) dt = \widehat{\psi}_\lambda(-Q)$ is

$$\begin{aligned} K_{\psi_\lambda}(x, y) &= \sum_{k=1}^N \chi_k(x) \iint \psi(t) e^{i(-t\lambda + \varphi_k(t;x,\xi) - y\xi)} a_k(t; x, \xi) dt d\xi \theta_k(y) \\ &\quad + \int e^{-it\lambda} \psi(t) r(t; x, y) dt, \end{aligned}$$

where the local coordinates in the right hand side depend on k and, to simplify the notation, we have omitted the corresponding coordinate maps. By the choices of ψ , θ_k and χ_k we obtain

$$\begin{aligned} (3.7) \quad \sum_j \widehat{\psi}(\lambda - \eta_j) &= \sum_{k=1}^N \iiint \psi(t) e^{i(-t\lambda + \varphi_k(t;x,\xi) - x\xi)} a_k(t; x, \xi) \theta_k(x) dt d\xi dx \\ &\quad + \iint e^{-it\lambda} \psi(t) r(t; x, x) dt dx \\ &= \sum_{k=1}^N \iiint \psi(t) e^{i(-t\lambda + \varphi_k(t;x,\xi) - x\xi)} a_k(t; x, \xi) \theta_k(x) dt d\xi dx \\ &\quad + O(|\lambda|^{-\infty}). \end{aligned}$$

³Trivially, for $j = 1, \dots, N-1$, q_k and a_k can be considered SG-classical, since, in those cases, they actually have order $-\infty$ with respect to x , by the fact that $q_k(x, \xi)$ vanishes for x outside a compact set.

Let $\psi \in C_0^\infty((-T, T))$, $T > 0$, be such that $\psi(0) = 1$ and $\widehat{\psi} \geq 0$, $\widehat{\psi}(0) > 0$ (e.g., set $\psi = \chi * \bar{\chi}$ with a suitable $\chi \in C_0^\infty((-T, T))$). By the analysis of the asymptotic behaviour of the integrals appearing in (3.7), described in Section 4, we finally obtain

$$(3.8) \quad \sum_j \widehat{\psi}(\lambda - \eta_j) = \begin{cases} \frac{n}{m} d_0 \lambda^{\frac{n}{m}-1} + O(\lambda^{n^*-1}) & \text{for } \lambda \rightarrow +\infty \\ O(|\lambda|^{-\infty}) & \text{for } \lambda \rightarrow -\infty, \end{cases}$$

with $n^* = \min\left\{n, \frac{n}{m} - 1\right\}$. The following Tauberian Theorem is a slight modification of Theorem 4.2.5 of [16] (see the Appendix):

Theorem 3.6. *Assume that*

- i) $\psi \in C_0^\infty(\mathbb{R})$ is an even function satisfying $\psi(0) = 1$, $\widehat{\psi} \geq 0$, $\widehat{\psi}(0) > 0$;
- ii) $N_Q(\lambda)$ is a nondecreasing function, supported in $[0, +\infty)$, continuous from the right, with polynomial growth at infinity and isolated discontinuity points of first kind $\{\eta_j\}$, $j \in \mathbb{N}$, such that $\eta_j \rightarrow +\infty$;
- iii) there exists $d_0 \geq 0$ such that

$$\sum_j \widehat{\psi}(\lambda - \eta_j) = \int \widehat{\psi}(\lambda - \eta) dN_Q(\eta) = \begin{cases} \frac{n}{m} d_0 \lambda^{\frac{n}{m}-1} + O(\lambda^{n^*-1}) & \text{for } \lambda \rightarrow +\infty \\ O(|\lambda|^{-\infty}) & \text{for } \lambda \rightarrow -\infty, \end{cases}$$

$$\text{with } m \in (0, 1), n^* = \min\left\{n, \frac{n}{m} - 1\right\}.$$

Then

$$N_Q(\lambda) = \frac{d_0}{2\pi} \lambda^{\frac{n}{m}} + O(\lambda^{n^*}), \text{ for } \lambda \rightarrow +\infty.$$

Remark 3.7. *The above statement can be modified as follows: with ψ , N_Q and m as in Theorem 3.6, when*

$$\int \widehat{\psi}(\lambda - \eta) dN_Q(\eta) = \begin{cases} \frac{n}{m} d_0 \lambda^{\frac{n}{m}-1} + O(\lambda^{\frac{n}{m}-2}) + O(\lambda^{n-1}) & \text{for } \lambda \rightarrow +\infty \\ O(|\lambda|^{-\infty}) & \text{for } \lambda \rightarrow -\infty, \end{cases}$$

$$\text{with } m \in (0, 1), \text{ then } N_Q(\lambda) = \frac{d_0}{2\pi} \lambda^{\frac{n}{m}} + O(\lambda^{\frac{n}{m}-1}) + O(\lambda^n), \text{ for } \lambda \rightarrow +\infty.$$

4. PROOF OF THEOREM 1.1

In view of Theorem 3.6 and Remark 3.7, to complete the proof of Theorem 1.1 we need to show that (3.8) holds. To this aim, as explained above, this Section will be devoted to studying the asymptotic behaviour for $|\lambda| \rightarrow +\infty$ of

$$(4.1) \quad I(\lambda) = \int e^{i\Phi(t; x, \xi; \lambda)} \psi(t) a(t; x, \xi) dt d\xi dx,$$

where $\psi \in C_0^\infty((-T, T))$, $\psi(0) = 1$, $a \in C^\infty((-T, T), S^{0,0}(\mathbb{R}^n))$, $a(0; x, \xi) = 1$, and

$$\Phi(t; x, \xi; \lambda) = \varphi(t; x, \xi) - x\xi - t\lambda, \quad \varphi \in C^\infty((-T, T), S_{\text{cl}}^{1,1}(\mathbb{R}^n))$$

such that

- $\partial_t \varphi(t; x, \xi) = q(x, d_x \varphi(t; x, \xi))$, $\varphi(0; x, \xi) = x\xi$;
- $C^{-1}\langle \xi \rangle \leq \langle d_x \varphi(t; x, \xi) \rangle \leq C\langle \xi \rangle$, for a suitable constant $C > 1$;
- $q \in S_{\text{cl}}^{m,1}(\mathbb{R}^n)$, $0 < m < 1$, SG-elliptic.

Since $q^{-1}(x, \xi) \in O(\langle x \rangle^{-1} \langle \xi \rangle^{-m})$ for $|x| + |\xi| \geq R > 0$, it is not restrictive to assume that this estimate holds⁴ on the whole phase space, so that, for a certain constant $A > 1$,

$$(4.2) \quad A^{-1} \langle x \rangle \langle \xi \rangle^m \leq q(x, \xi) \leq A \langle x \rangle \langle \xi \rangle^m.$$

For two functions f, g , defined on a common subset X of \mathbb{R}^{d_1} and depending on parameters $y \in Y \subseteq \mathbb{R}^{d_2}$, we will write $f < g$ or $f(x, y) < g(x, y)$ to mean that there exists a suitable constant $c > 0$ such that $|f(x, y)| \leq c|g(x, y)|$ for all $(x, y) \in X \times Y$. The notation $f \sim g$ or $f(x, y) \sim g(x, y)$ means that both $f < g$ and $g < f$ hold.

Remark 4.1. *The ellipticity of q yields, for $\lambda < 0$,*

$$\partial_t \Phi(t; x, \xi; \lambda) = q(x, d_x \varphi(t; x, \xi)) - \lambda > \langle x \rangle \langle \xi \rangle^m + |\lambda|$$

which, by integration by parts, implies $I(\lambda) = O(|\lambda|^{-\infty})$ when $\lambda \rightarrow -\infty$.

From now on any asymptotic estimate is to be meant for $\lambda \rightarrow +\infty$.

We will make use of a partition of unity on the phase space: the supports of its elements will depend on suitably large positive constants $k_1, k_2 > 1$. We also assume, as it is possible, $\lambda \geq \lambda_0$, again with an appropriate $\lambda_0 \gg 1$. As we will see below, the values of k_1, k_2 and λ_0 depend only on q and its associated seminorms.

Proposition 4.2. *Let H_1 be any function in $C_0^\infty(\mathbb{R})$ such that $\text{supp } H_1 \subseteq [(2k_1)^{-1}, 2k_1]$, $0 \leq H_1 \leq 1$ and $H_1 \equiv 1$ on $[k_1^{-1}, k_1]$, where $k_1 > 1$ is a suitably chosen, large positive constant. Then*

$$(4.3) \quad I(\lambda) = O(\lambda^{-\infty}) + \int e^{i\Phi(t; x, \xi; \lambda)} \psi(t) H_1\left(\frac{\langle x \rangle \langle \xi \rangle^m}{\lambda}\right) a(t; x, \xi) dt d\xi dx.$$

Proof. Write

$$(4.4) \quad \begin{aligned} I(\lambda) = & \int e^{i\Phi(t; x, \xi; \lambda)} \psi(t) \left[1 - H_1\left(\frac{\langle x \rangle \langle \xi \rangle^m}{\lambda}\right) \right] a(t; x, \xi) dt d\xi dx \\ & + \int e^{i\Phi(t; x, \xi; \lambda)} \psi(t) H_1\left(\frac{\langle x \rangle \langle \xi \rangle^m}{\lambda}\right) a(t; x, \xi) dt d\xi dx \end{aligned}$$

and observe that, by $A^{-1} \langle x \rangle \langle \xi \rangle^m \leq q(x, \xi) \leq A \langle x \rangle \langle \xi \rangle^m$, $x, \xi \in \mathbb{R}^n$, we find

$$\begin{aligned} |\partial_t \Phi(t; x, \xi; \lambda)| &\geq \frac{\lambda}{2} + \left(\frac{k_1}{2} - AC \right) \langle x \rangle \langle \xi \rangle^m && \text{when } \frac{\langle x \rangle \langle \xi \rangle^m}{\lambda} \leq k_1^{-1}, \\ |\partial_t \Phi(t; x, \xi; \lambda)| &\geq \frac{(AC)^{-1}}{2} \langle x \rangle \langle \xi \rangle^m + \left[\frac{(AC)^{-1}}{2} k_1 - 1 \right] \lambda && \text{when } \frac{\langle x \rangle \langle \xi \rangle^m}{\lambda} \geq k_1. \end{aligned}$$

Thus, if $k_1 > 2AC$ we have $|\partial_t \Phi(t; x, \xi; \lambda)| \sim \lambda + \langle x \rangle \langle \xi \rangle^m$ on the support of $1 - H_1\left(\frac{\langle x \rangle \langle \xi \rangle^m}{\lambda}\right)$, and the assertion follows integrating by parts with respect to t in the first integral of (4.4). \square

Remark 4.3. *We actually choose $k_1 > 4AC > 2AC$, since this will be needed in the proof of Proposition 4.7 below, see also subsection A.3 in the Appendix.*

⁴This amounts, at most, to modifying q by adding and subtracting a compactly supported symbol, that is, an element of $S^{-\infty, -\infty}(\mathbb{R}^n)$. The corresponding solutions φ and a of the eikonal and transport equations, respectively, would then change, at most, by an element of $C^\infty((-T, T), S^{-\infty, -\infty}(\mathbb{R}^n))$, see [8, 10, 11]: it is immediate, by integration by parts with respect to t , that an integral as (4.1) is $O(|\lambda|^{-\infty})$ for $a \in C^\infty((-T, T), S^{-\infty, -\infty}(\mathbb{R}^n))$. Then, the modified q obviously keeps the same sign everywhere.

Let us now pick $H_2 \in C_0^\infty(\mathbb{R})$ such that $0 \leq H_2(v) \leq 1$, $H_2(v) = 1$ for $|v| \leq k_2$ and $H_2(v) = 0$ for $|v| \geq 2k_2$, where $k_2 > 1$ is a constant which we will choose big enough (see below). We can then write

$$\begin{aligned} I(\lambda) &= O(\lambda^{-\infty}) + \int e^{i\Phi(t;x,\xi;\lambda)} \psi(t) H_1\left(\frac{\langle x \rangle \langle \xi \rangle^m}{\lambda}\right) H_2(|\xi|) a(t; x, \xi) dt d\xi dx \\ &\quad + \int e^{i\Phi(t;x,\xi;\lambda)} \psi(t) H_1\left(\frac{\langle x \rangle \langle \xi \rangle^m}{\lambda}\right) [1 - H_2(|\xi|)] a(t; x, \xi) dt d\xi dx \\ &= O(\lambda^{-\infty}) + I_1(\lambda) + I_2(\lambda). \end{aligned}$$

In what follows, we will sistematically use the notation $S^{r,\rho} = S^{r,\rho}(y, \eta)$, $y \in \mathbb{R}^k$, $\eta \in \mathbb{R}^l$, to generally denote functions depending smoothly on y and η and satisfying SG-type estimates of order r, ρ in y, η . In a similar fashion, $S_T^{r,\rho} = C^\infty((-T, T), S^{r,\rho}(y, \eta))$ will stand for some function of the same kind which, additionally, depends smoothly on $t \in (-T, T)$, and, for all $s \in \mathbb{Z}_+$, $D_t^s C^\infty((-T, T), S^{r,\rho}(y, \eta))$ satisfies SG-type estimates of order r, ρ in y, η , uniformly with respect to $t \in (-T, T)$.

To estimate $I_1(\lambda)$, we will apply the Stationary Phase Theorem. We begin by rewriting the integral $I_1(\lambda)$, using the fact that φ is solution of the eikonal equation associated with q and that q is a classical SG-symbol. Note that then $\partial_t^2 \varphi \in C^\infty((-T, T), S_{\text{cl}}^{2m-1,1}(\mathbb{R}^n)) \subseteq C^\infty((-T, T), S_{\text{cl}}^{m,1}(\mathbb{R}^n))$, since

$$\partial_t^2 \varphi(t; x, \xi) = \sum_{i=1}^n (\partial_{\xi_i} q)(x, d_x \varphi(t; x, \xi)) \partial_{x_i} (q(x, d_x \varphi(t; x, \xi))).$$

In view of the Taylor expansion of φ at $t = 0$, recalling the property $q(x, \xi) = \omega(x)q_e(x, \xi) + S^{m,0}(x, \xi)$, ω a fixed 0-excision function, we have, for some $0 < \delta_1 < 1$,

$$\begin{aligned} \Phi(t; x, \xi; \lambda) &= -\lambda t - x\xi + \varphi(0; x, \xi) + t \partial_t \varphi(0; x, \xi) + \frac{t^2}{2} \partial_t^2 \varphi(t\delta_1; x, \xi) \\ &= -\lambda t + tq(x, \xi) + t^2 S_T^{2m-1,1}(x, \xi) \\ &= -\lambda t + t\omega(x)q_e(x, \xi) + tS^{m,0}(x, \xi) + t^2 S_T^{2m-1,1}(x, \xi) \\ &= -\lambda t + t\omega(x)q_e(x, \xi) + tS^{m,0}(x, \xi) + t^2 \omega(x)S_{T,e}^{2m-1,1}(x, \xi) + t^2 S_T^{2m-1,0}(x, \xi), \end{aligned}$$

where the subscript e denotes the x -homogeneous (exit) principal parts of the involved symbols, which are all SG-classical and real-valued, see [10].

Observe that $|x| \sim \lambda$ on the support of the integrand in $I_1(\lambda)$, so that we can, in fact, assume⁵ $\omega(x) \equiv 1$ there. Indeed,

$$|\xi| < 1, \langle x \rangle \langle \xi \rangle^m \sim \lambda \Rightarrow \langle x \rangle \sim \lambda,$$

which of course implies $\langle x \rangle \sim |x|$, provided $\lambda_0 \leq \lambda$ is large enough. Moreover, by the ellipticity of q , writing $x = |x|\varsigma$, $\varsigma \in \mathbb{S}^{n-1}$, with the constant $A > 1$ of (4.2),

$$\begin{aligned} A^{-1} \langle x \rangle \langle \xi \rangle^m &\leq q(x, \xi) = \omega(x)q_e(x, \xi) + S^{m,0}(x, \xi) \leq A \langle x \rangle \langle \xi \rangle^m \\ &\Rightarrow A^{-1} \frac{\langle x \rangle}{|x|} \langle \xi \rangle^m \leq \omega(x)q_e(\varsigma, \xi) + \frac{S^{m,0}(x, \xi)}{|x|} \leq A \frac{\langle x \rangle}{|x|} \langle \xi \rangle^m \\ (4.5) \quad &\Rightarrow A^{-1} \langle \xi \rangle^m \leq q_e(\varsigma, \xi) \leq A \langle \xi \rangle^m, \quad \varsigma \in \mathbb{S}^{n-1}, \xi \in \mathbb{R}^n \end{aligned}$$

⁵By definition, $\omega \in C^\infty(\mathbb{R}^n)$, $\omega(v) \equiv 0$ for $|v| \leq B$, $\omega(v) \equiv 1$ for $|v| \geq 2B$, for a constant $B > 0$.

taking the limit for $|x| \rightarrow +\infty$. Then, setting $x = \lambda\zeta\varsigma$, $\zeta \in [0, +\infty)$, $\varsigma \in \mathbb{S}^{n-1}$, $\lambda \geq \lambda_0 \gg 1$, in $I_1(\lambda)$, by homogeneity and the above remarks, we can write

$$\begin{aligned} \Phi(t; \lambda\zeta\varsigma, \xi; \lambda) &= -\lambda t + t\omega(\lambda\zeta\varsigma)q_e(\lambda\zeta\varsigma, \xi) + tS^{m,0}(\lambda\zeta\varsigma, \xi) \\ &\quad + t^2\omega(\lambda\zeta\varsigma)S_{T,e}^{2m-1,1}(\lambda\zeta\varsigma, \xi) + t^2S_T^{2m-1,0}(\lambda\zeta\varsigma, \xi) \\ &= -\lambda t + \lambda\zeta t q_e(\varsigma, \xi) + \lambda\zeta t^2 S_{T,e}^{2m-1,1}(\varsigma, \xi) + tS^{m,0}(\lambda\zeta\varsigma, \xi) + t^2S_T^{2m-1,0}(\lambda\zeta\varsigma, \xi) \\ &= \lambda[-t + \zeta t q_e(\varsigma, \xi) + \zeta t^2 S_{T,e}^{2m-1,1}(\varsigma, \xi)] + G_1(\lambda; t, \zeta; \varsigma, \xi) \\ &= \lambda F_1(t, \zeta; \varsigma, \xi) + G_1(\lambda; t, \zeta; \varsigma, \xi), \end{aligned}$$

and find, in view of the compactness of the support of the integrand (see the proof of Proposition 4.4 below) and the hypotheses,

$$\begin{aligned} I_1(\lambda) &= \lambda^n \int e^{i\lambda F_1(t, \zeta; \varsigma, \xi)} e^{iG_1(\lambda; t, \zeta; \varsigma, \xi)} \psi(t) a(t; \lambda\zeta\varsigma, \xi) H_1\left(\frac{\langle \lambda\zeta \rangle \langle \xi \rangle^m}{\lambda}\right) H_2(|\xi|) \zeta^{n-1} dt d\zeta d\xi d\varsigma \\ &= \frac{\lambda^n}{(2\pi)^n} \int e^{i\lambda F_1(X, Y)} U_1(X, Y; \lambda) dX dY, \end{aligned}$$

with $X = (t, \zeta)$, $Y = (\varsigma, \xi)$. We can now prove

Proposition 4.4. *Choosing the constants $k_1, \lambda_0 > 1$ large enough and $T > 0$ suitably small, we have, for any $k_2 > 1$ and for a certain sequence c_j , $j = 0, 1, \dots$,*

$$I_1(\lambda) \sim \sum_{j=0}^{+\infty} c_j \lambda^{n-1-j},$$

that is, $I_1(\lambda) = c_0 \lambda^{n-1} + O(\lambda^{n-2})$, with

$$c_0 = \frac{1}{(2\pi)^{n-1}} \int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} \frac{H_2(|\xi|)}{q_e(\varsigma, \xi)^n} d\varsigma d\xi.$$

Proof. It is easy to see that, on the support of U_1 , the phase function $F_1(X, Y)$ admits a unique, nondegenerate, stationary point $X_0 = X_0(Y) = (0, q_e(\varsigma, \xi)^{-1})$, i.e. $F'_{1,X}(X_0(Y), Y) = 0$ for all Y such that $(X, Y) \in \text{supp } U_1$, provided $T > 0$ is chosen suitably small (see, e.g., [14], p. 136), and the Hessian $\det(F''_{1,X}(X_0(Y), Y))$ equals $-q_e(\varsigma, \xi)^2 < 0$. Moreover, the amplitude function

$$U_1(X, Y; \lambda) = \psi(t) H_1\left(\frac{\langle \lambda\zeta \rangle \langle \xi \rangle^m}{\lambda}\right) H_2(|\xi|) a(t; \lambda\zeta\varsigma, \xi) \zeta^{n-1} e^{iG(\lambda; t, \zeta; \varsigma, \xi)}$$

is compactly supported with respect to the variables X and Y , and satisfies, for all $\gamma \in \mathbb{Z}_+^2$,

$$D_X^\gamma U_1(X, Y; \lambda) < 1$$

for all X, Y , $\lambda \geq \lambda_0$. In fact:

- (1) $\psi \in C_0^\infty((-T, T))$, $\varsigma \in \mathbb{S}^{n-1}$, $\text{supp}[H_2(|\xi|)] \subseteq \{\xi: |\xi| \leq 2k_2\}$, and

$$(2k_1)^{-1} \leq \langle \xi \rangle^m \sqrt{\frac{1}{\lambda^2} + \zeta^2} \leq 2k_1 \Rightarrow 0 < \sqrt{\frac{1}{4k_1^2 \langle 2k_2 \rangle^{2m}} - \frac{1}{\lambda_0^2}} \leq \zeta \leq 2k_1,$$

where $\lambda_0 > 2k_1 \langle 2k_2 \rangle^m$;

- (2) all the factors appearing in the expression of U_1 are uniformly bounded, together with all their X -derivatives, for $X \in S_X = \text{supp } \psi \times [\zeta_0, \zeta_1]$, $Y \in S_Y = \mathbb{S}^{n-1} \times \{\xi: |\xi| \leq 2k_2\}$, and $\lambda \geq \lambda_0$.

Of course, (2) trivially holds for the cutoff functions $\psi(t)$ and $H_2(|\xi|)$, and for the factor ζ^{n-1} . Since $a(t; x, \xi) \in S_T^{0,0}(x, \xi)$, on $S_X \times S_Y$ we have, for all $\gamma \in \mathbb{Z}_+^2$ and $\lambda \geq \lambda_0 > 1$,

$$D_X^\gamma a(t; \lambda \zeta, \xi) < \langle \lambda \zeta \rangle^{-\gamma_2} \lambda^{\gamma_2} \langle \xi \rangle^m < \frac{1}{\left(\frac{1}{\lambda^2} + \zeta^2\right)^{\frac{\gamma_2}{2}}} < \frac{1}{\zeta^{\gamma_2}} < 1.$$

Moreover, since $G_1 \in S_T^{m,0}(x, \xi)$ is actually in $S_T^{-\infty,0}(x, \xi) \subset S_T^{0,0}(x, \xi)$ on $S_X \times S_Y$, the same holds for $\exp(iG_1)$, by an application of the Faà di Bruno formula for the derivatives of compositions of functions, so also this factor fulfills the desired estimates. Finally, another straightforward computation shows that, for all $\gamma_2 \in \mathbb{Z}_+$ and $\lambda \geq \lambda_0 > 1$,

$$D_\zeta^{\gamma_2} H_1\left(\frac{\langle \lambda \zeta \rangle \langle \xi \rangle^m}{\lambda}\right) < 1$$

on $S_X \times S_Y$. The Proposition is then a consequence of the Stationary Phase Theorem (see [12], Proposition 1.2.4, [21], Theorem 7.7.6), applied to the integral with respect to $X = (t, \zeta)$. In particular, the leading term is given by $\frac{\lambda^n}{(2\pi)^{n-1}}$ times the integral with respect to Y of $\lambda^{-1} |\det(F''_{1,X}(X_0(Y), Y))|^{-\frac{1}{2}} U_1(X_0(Y), Y; \lambda)$, that is⁶

$$\begin{aligned} I_1(\lambda) &= \frac{\lambda^{n-1}}{(2\pi)^{n-1}} \int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} \frac{1}{q_e(\zeta, \xi)} \psi(0) H_1\left(\frac{\langle \frac{\lambda}{q_e(\zeta, \xi)} \rangle \langle \xi \rangle^m}{\lambda}\right) \frac{H_2(|\xi|)}{q_e(\zeta, \xi)^{n-1}} a\left(0; \frac{\lambda \zeta}{q_e(\zeta, \xi)}, \xi\right) d\zeta d\xi \\ &\quad + O(\lambda^{n-2}) \\ &= \frac{\lambda^{n-1}}{(2\pi)^{n-1}} \int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} H_1\left(\frac{\langle \frac{\lambda}{q_e(\zeta, \xi)} \rangle \langle \xi \rangle^m}{\lambda}\right) \frac{H_2(|\xi|)}{q_e(\zeta, \xi)^n} d\zeta d\xi + O(\lambda^{n-2}) \\ &= \frac{\lambda^{n-1}}{(2\pi)^{n-1}} \int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} \frac{H_2(|\xi|)}{q_e(\zeta, \xi)^n} d\zeta d\xi + O(\lambda^{n-2}). \end{aligned}$$

Indeed, having chosen $k_1 > 2A$, $\lambda_0 > 2k_1 \langle 2k_2 \rangle^m$, (4.5) implies

$$k_1^{-1} < A^{-1} < \frac{\langle \frac{\lambda}{q_e(\zeta, \xi)} \rangle \langle \xi \rangle^m}{\lambda} = \sqrt{\left(\frac{\langle \xi \rangle^m}{\lambda}\right)^2 + \left(\frac{\langle \xi \rangle^m}{q_e(\zeta, \xi)}\right)^2} < \sqrt{\frac{1}{4k_1^2} + A^2} < k_1,$$

uniformly for $\zeta \in \mathbb{S}^{n-1}$, $\xi \in \text{supp}[H_2(|\xi|)]$, $\lambda \geq \lambda_0$. This concludes the proof. \square

Let us now consider $I_2(\lambda)$. We follow a procedure close to that used in the proof of Theorem 7.7.6 of [21]. However, since here we lack the compactness of the support of the amplitude with respect to x , we need explicit estimates to show that all the involved integrals are convergent, so we give below the argument in full detail.

We initially proceed as in the analysis of $I_1(\lambda)$ above. In view of the presence of the factor $1 - H_2(|\xi|)$ in the integrand, we can now assume $|\xi| \geq k_2 > \max\{B, 1\}$, $B > 0$ the radius of the smallest ball in \mathbb{R}^n including $\text{supp}(1 - \omega)$, so that $q(x, \xi) = \omega(\xi)q_\psi(x, \xi) + S^{m-1,1}(x, \xi) = q_\psi(x, \xi) + S^{m-1,1}(x, \xi)$. Then, with some $0 < \delta_2 < 1$,

$$\begin{aligned} \Phi(t; x, \xi; \lambda) &= -\lambda t - x\xi + \varphi(0; x, \xi) + t \partial_t \varphi(0; x, \xi) + \frac{t^2}{2} \partial_t^2 \varphi(t\delta_2; x, \xi) \\ &= -\lambda t + tq(x, \xi) + t^2 S_T^{2m-1,1}(x, \xi) \\ &= -\lambda t + tq_\psi(x, \xi) + t S^{m-1,1}(x, \xi) + t^2 S_T^{2m-1,1}(x, \xi). \end{aligned}$$

⁶Remember that $\psi(0) = 1$, $a(0; x, \xi) = 1$ for all $x, \xi \in \mathbb{R}^n$.

Setting $\xi = (\lambda\zeta)^{\frac{1}{m}}\varsigma$, $\zeta \in [0, +\infty)$, $\varsigma \in \mathbb{S}^{n-1}$, $\lambda \geq \lambda_0$, we can rewrite $I_2(\lambda)$ as

$$\begin{aligned} I_2(\lambda) &= \frac{n}{m} \frac{\lambda^{\frac{n}{m}}}{(2\pi)^n} \int e^{i\lambda(-t+\zeta tq_\psi(x,\varsigma)+t\lambda^{-1}S^{m-1,1}(x,(\lambda\zeta)^{\frac{1}{m}}\varsigma)+t^2\lambda^{-1}S_T^{2m-1,1}(x,(\lambda\zeta)^{\frac{1}{m}}\varsigma))} \\ &\quad \cdot \psi(t) a(t; x, (\lambda\zeta)^{\frac{1}{m}}\varsigma) H_1\left(\frac{\langle x \rangle \langle (\lambda\zeta)^{\frac{1}{m}}\varsigma \rangle^m}{\lambda}\right) \left[1 - H_2((\lambda\zeta)^{\frac{1}{m}}\varsigma)\right] \zeta^{\frac{n}{m}-1} dt d\zeta d\varsigma dx \\ &= \frac{n}{m} \frac{\lambda^{\frac{n}{m}}}{(2\pi)^n} \int e^{i\lambda F_2(X,Y;\lambda)} U_2(X,Y;\lambda) dXdY, \end{aligned}$$

$X = (t, \zeta)$, $Y = (\varsigma, x)$, where we have set

$$F_2(X,Y;\lambda) = -t + \zeta tq_\psi(x,\varsigma) + t\lambda^{-1}S^{m-1,1}(x,(\lambda\zeta)^{\frac{1}{m}}\varsigma) + t^2\lambda^{-1}S_T^{2m-1,1}(x,(\lambda\zeta)^{\frac{1}{m}}\varsigma)$$

$$U_2(X,Y;\lambda) = \psi(t) H_1\left(\frac{\langle x \rangle \langle (\lambda\zeta)^{\frac{1}{m}}\varsigma \rangle^m}{\lambda}\right) \left[1 - H_2((\lambda\zeta)^{\frac{1}{m}}\varsigma)\right] a(t; x, (\lambda\zeta)^{\frac{1}{m}}\varsigma) \zeta^{\frac{n}{m}-1}.$$

On the support of U_2 we have

$$\frac{\langle x \rangle \langle (\lambda\zeta)^{\frac{1}{m}}\varsigma \rangle^m}{\lambda} \sim 1 \quad \text{and} \quad (\lambda\zeta)^{\frac{1}{m}} > 1 \Rightarrow \langle (\lambda\zeta)^{\frac{1}{m}}\varsigma \rangle^m = \langle (\lambda\zeta)^{\frac{1}{m}} \rangle^m \sim \lambda\zeta,$$

so that

$$(4.6) \quad \frac{\langle x \rangle \lambda \zeta}{\lambda} \sim 1 \Leftrightarrow \zeta \sim \langle x \rangle^{-1} \quad \text{and} \quad |x| < \langle x \rangle \leq 2k_1 \langle k_2 \rangle^{-m} \lambda = \tilde{\kappa} \lambda.$$

For any fixed $Y \in \mathbb{S}^{n-1} \times \mathbb{R}^n$ we then have X belonging to a compact set, uniformly with respect to $\lambda \geq \lambda_0$, say $\text{supp } \psi \times [c^{-1}\langle x \rangle^{-1}, c\langle x \rangle^{-1}]$, for a suitable $c > 1$.

Remark 4.5. Incidentally, we observe that a rough estimate of $\lambda^{\frac{n}{m}} I_2(\lambda)$ is

$$\begin{aligned} \int e^{i\lambda F_2(X,Y;\lambda)} U_2(X,Y;\lambda) dX &< \langle x \rangle^{-\frac{n}{m}+1} \int_{c^{-1}\langle x \rangle^{-1}}^{c\langle x \rangle^{-1}} d\zeta < \langle x \rangle^{-\frac{n}{m}} \\ &\Rightarrow \lambda^{\frac{n}{m}} \int e^{i\lambda F_2(X,Y;\lambda)} U_2(X,Y;\lambda) dXdY < \lambda^n, \lambda \rightarrow +\infty. \end{aligned}$$

An even less precise result would be the bound $\lambda^{\frac{n}{m}}$, using the convergence of the integral with respect to x in the whole \mathbb{R}^n , given by $-\frac{n}{m} + n < 0$.

The next Lemma is immediate, and we omit the proof:

Lemma 4.6. $S_T^{s,\sigma}(x, (\lambda\zeta)^{\frac{1}{m}}\varsigma) = S_T^{s,\sigma}(x, (\lambda\zeta)^{\frac{1}{m}}\varsigma)$ for any $\zeta \in [0, +\infty)$, $x \in \mathbb{R}^n$, $\varsigma \in \mathbb{S}^{n-1}$, $\lambda \geq \lambda_0$, $m \in (0, 1)$, and, for all $\gamma \in \mathbb{Z}_+^2$,

$$D_X^\gamma S_T^{s,\sigma}(x, (\lambda\zeta)^{\frac{1}{m}}\varsigma) = \zeta^{-\gamma_2} S_T^{s,\sigma}(x, (\lambda\zeta)^{\frac{1}{m}}\varsigma).$$

The main result of this Section is

Proposition 4.7. If $k_1, k_2, \lambda_0 > 1$ are chosen large enough we have

$$(4.7) \quad I_2(\lambda) = \frac{n}{m} d_0 \lambda^{\frac{n}{m}-1} + O(\lambda^{n-1}) + O(\lambda^{\frac{n}{m}-2}).$$

Explicitly,

$$d_0 = \frac{1}{(2\pi)^{n-1}} \int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} \frac{1}{q_\psi(x, \varsigma)^{\frac{n}{m}}} d\varsigma dx.$$

We will prove Proposition 4.7 through various intermediate steps. First of all, arguing as in the proof of (4.5), exchanging the role of x and ξ , we note that, for all $x \in \mathbb{R}^n$, $\varsigma \in \mathbb{S}^{n-1}$,

$$(4.8) \quad A^{-1}\langle x \rangle \leq q_\psi(x, \varsigma) \leq A\langle x \rangle,$$

$(x, \varsigma) \in \mathbb{R}^n \times \mathbb{S}^{n-1}$. We now study

$$\begin{aligned} F'_{2,X}(X, Y; \lambda) &= \begin{pmatrix} \partial_t F_2(X, Y; \lambda) \\ \partial_\zeta F_2(X, Y; \lambda) \end{pmatrix} \\ &= \begin{pmatrix} -1 + \frac{\zeta}{\zeta_0} + \lambda^{-1} S^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) + t \lambda^{-1} S_T^{2m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) \\ t(q_\psi(x, \varsigma) + \lambda^{-1} \zeta^{-1} S^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) + t \lambda^{-1} \zeta^{-1} S_T^{2m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}})) \end{pmatrix}, \end{aligned}$$

$X = (t, \zeta) \in S_X = \text{supp } \psi \times [c^{-1}\langle x \rangle^{-1}, c\langle x \rangle^{-1}]$, $Y = (\varsigma, x) \in S_Y = \mathbb{S}^{n-1} \times \mathbb{R}^n$, $\lambda \geq \lambda_0$, where we have used Lemma 4.6. By the symbolic calculus, remembering that $\lambda \zeta \geq k_2^m > 1$ on $\text{supp } U_2$, we can rewrite the expressions above as

$$\begin{aligned} \partial_t F_2(X, Y; \lambda) &= -1 + \frac{\zeta}{\zeta_0} + \zeta(\lambda \zeta)^{-1} S^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) + t \zeta(\lambda \zeta)^{-1} S_T^{2m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) \\ &= -1 + \frac{\zeta}{\zeta_0} + \zeta[(\lambda \zeta)^{\frac{1}{m}}]^{-m} S^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) + t \zeta[(\lambda \zeta)^{\frac{1}{m}}]^{-m} S_T^{2m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) \\ &= -1 + \frac{\zeta}{\zeta_0} + \zeta S^{-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) + t \zeta S_T^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}), \\ \partial_\zeta F_2(X, Y; \lambda) &= t(q_\psi(x, \varsigma) + S^{-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) + t S_T^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}})). \end{aligned}$$

It is clear that $\zeta \sim \langle x \rangle^{-1}$ implies $\zeta S^{-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) = S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}})$ and $\zeta S_T^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) = S_T^{m-1,0}(x, (\lambda \zeta)^{\frac{1}{m}})$, so that we finally have

$$\begin{aligned} \partial_t F_2(X, Y; \lambda) &= -1 + \frac{\zeta}{\zeta_0} + S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}}) + t S_T^{m-1,0}(x, (\lambda \zeta)^{\frac{1}{m}}), \\ \partial_\zeta F_2(X, Y; \lambda) &= t(q_\psi(x, \varsigma) + S^{-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) + t S_T^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}})). \end{aligned}$$

We now prove that, modulo an $O(|\lambda|^{-\infty})$ term, we can consider an amplitude such that, on its support, the ration ζ/ζ_0 is very close to 1. To this aim, take $H_3 \in C_0^\infty(\mathbb{R})$ such that $0 \leq H_3(v) \leq 1$, $H_3(v) = 1$ for $|v| \leq \frac{3}{2}\varepsilon$ and $H_3(v) = 0$ for $|v| \geq 2\varepsilon$, with an arbitrarily fixed, small enough $\varepsilon \in (0, \frac{1}{2})$, and set

$$V_1(X, Y; \lambda) = U_2(X, Y; \lambda) \cdot \left[1 - H_3\left(\frac{\zeta}{\zeta_0} - 1\right) \right], \quad V_2(X, Y; \lambda) = U_2(X, Y; \lambda) \cdot H_3\left(\frac{\zeta}{\zeta_0} - 1\right),$$

$$J_1(\lambda) = \int e^{i\lambda F_2(X, Y; \lambda)} V_1(X, Y; \lambda) dX dY, \quad J_2(\lambda) = \int e^{i\lambda F_2(X, Y; \lambda)} V_2(X, Y; \lambda) dX dY.$$

Proposition 4.8. *With the choices of T, k_1, λ_0 above, for any $\varepsilon \in (0, \frac{1}{2})$ we can find $k_2 > 1$ large enough such that $J_1(\lambda) = O(\lambda^{-\infty})$.*

Proof. Since $0 < m < 1$, in view of (1.1), (4.6), and (4.8), we can choose $k_2 > 1$ so large that, for an arbitrarily fixed $\varepsilon \in (0, \frac{1}{2})$, for any $\lambda \geq \lambda_0$, $\zeta \in (0, +\infty)$ satisfying $|\xi| = (\lambda \zeta)^{\frac{1}{m}} \geq k_2$,

$$\begin{aligned} (4.9) \quad & \text{in } \partial_t F_2(X, Y; \lambda), \quad \left| S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}}) \right| \leq \frac{\varepsilon}{2}, \quad \left| t S_T^{m-1,0}(x, (\lambda \zeta)^{\frac{1}{m}}) \right| \leq \frac{\varepsilon}{2}, \\ & \text{and } \left| \zeta_0 \frac{d}{d\zeta} S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}}) \right| = \left| \zeta_0 \zeta^{-1} S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}}) \right| \leq k_0 < 1, \\ & \text{in } \partial_\zeta F_2(X, Y; \lambda), \quad |S^{-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) + t S_T^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}})| \leq \frac{A^{-1}}{2} \langle x \rangle, \end{aligned}$$

uniformly with respect to $(X, Y) \in S_X \times S_Y \supseteq \text{supp } U_2(\cdot; \lambda)$. Then, F_2 is non-stationary on $\text{supp } V_1$, since there we have $\left| \frac{\zeta}{\zeta_0} - 1 \right| \geq \frac{3}{2}\varepsilon$, while

$$\left| S^{-1,0}(x, (\lambda\zeta)^{\frac{1}{m}}) + tS_T^{m-1,0}(x, (\lambda\zeta)^{\frac{1}{m}}) \right| \leq \varepsilon,$$

which implies $\partial_t F_2(X, Y; \lambda) > 1$. Observing that, on $\text{supp } V_1$, $\partial_t F_2(X, Y; \lambda) = S_T^{0,0}(x, (\lambda\zeta)^{\frac{1}{m}})$, as well as $V_1(X, Y; \lambda) = S_T^{0,0}(x, (\lambda\zeta)^{\frac{1}{m}})$, the assertion follows by repeated integrations by parts with respect to t , using the operator

$$L_t = \frac{1}{\lambda \partial_t F_2(X, Y; \lambda)} D_t \Rightarrow L_1 e^{i\lambda F_2(X, Y; \lambda)} = e^{i\lambda F_2(X, Y; \lambda)}$$

and recalling Remark 4.5. \square

Proposition 4.9. *With the choices of $\varepsilon, T > 0, k_1, k_2, \lambda_0 > 1$ above, we can assume, modulo an $O(\lambda^{n-1})$ term, that the integral with respect to x in $J_2(\lambda)$ is extended to the set $\{x \in \mathbb{R}^n : \langle x \rangle \leq \kappa\lambda\}$, with*

$$(4.10) \quad \kappa = \left(1 - \frac{\varepsilon}{2}\right) [A(2k_2)^m]^{-1}.$$

Proof. Indeed⁷, if $\kappa < \tilde{\kappa} = 2k_1 \langle k_2 \rangle^{-m}$, we can split $J_2(\lambda)$ into the sum

$$(4.11) \quad \int_{\kappa\lambda \leq \langle x \rangle \leq \tilde{\kappa}\lambda} \int_{\mathbb{S}^{n-1}} \int e^{i\lambda F_2} V_2 dX d\zeta dx + \int_{\langle x \rangle \leq \kappa\lambda} \int_{\mathbb{S}^{n-1}} \int e^{i\lambda F_2} V_2 dX d\zeta dx.$$

Observing that, on $\text{supp } U_2$,

$$\langle x \rangle \sim \lambda \Rightarrow \langle \xi \rangle^m = \frac{\langle x \rangle \langle \xi \rangle^m}{\lambda} \frac{\lambda}{\langle x \rangle} \sim 1 \Rightarrow |\xi| \leq k_3,$$

switching back to the original variables, the first integral in (4.11) can be treated as $I_1(\lambda)$, and gives, in view of Proposition 4.4, an $O(\lambda^{n-1})$ term, as stated. \square

Now we can show that $F_2(X, Y; \lambda)$ admits a unique, nondegenerate stationary point $X_0^* = X_0^*(Y, \lambda)$ belonging to $\text{supp } V_2$ for $\langle x \rangle \leq \kappa\lambda$. Under the same hypotheses, X_0^* lies in a circular neighbourhood of $X_0 = (0, \zeta_0) = (0, q_\psi(x, \varsigma)^{-1})$ of arbitrarily small radius:

Proposition 4.10. *With $\varepsilon \in (0, \frac{1}{2})$, $T > 0, k_1, k_2, \lambda_0 > 1$ fixed above, $F'_{2,X}(X, Y; \lambda)$ vanishes on $\text{supp } V_2$ only for $X = X_0^*(Y; \lambda) = (0, \zeta_0^*(Y; \lambda))$, i.e., $F'_{2,X}(X_0^*(Y; \lambda), Y; \lambda) = 0$ for all Y such that $(X, Y; \lambda) \in \text{supp } V_2$. Moreover,*

$$\det(F''_{2,X}(X_0^*(Y; \lambda), Y)) \sim \langle x \rangle^2 \text{ and } |X_0^*(Y; \lambda) - X_0(Y)| = |\zeta_0^*(Y; \lambda) - \zeta_0(Y)| \leq \frac{A\varepsilon}{2} \langle x \rangle^{-1}$$

holds on $\text{supp } V_2$.

Proof. We have to solve

$$\begin{cases} 0 = -1 + \frac{\zeta}{\zeta_0} + S^{-1,0}(x, (\lambda\zeta)^{\frac{1}{m}}) + tS_T^{m-1,0}(x, (\lambda\zeta)^{\frac{1}{m}}) \\ 0 = t(q_\psi(x, \varsigma) + S^{-1,1}(x, (\lambda\zeta)^{\frac{1}{m}}) + tS_T^{m-1,1}(x, (\lambda\zeta)^{\frac{1}{m}})), \end{cases}$$

$(X, Y; \lambda) \in \text{supp } V_2$. By (4.8) and (4.9), with the choices of $\varepsilon, T > 0, k_1, k_2, \lambda_0$ above, the coefficient of t in the second equation does not vanish at any point of $\text{supp } V_2$. Then $t = 0$, and ζ must satisfy

$$(4.12) \quad -1 + \frac{\zeta}{\zeta_0} + S^{-1,0}(x, (\lambda\zeta)^{\frac{1}{m}}) = 0 \Leftrightarrow \zeta = \zeta_0(1 + S^{-1,0}(x, (\lambda\zeta)^{\frac{1}{m}})) = G(\zeta; Y; \lambda).$$

⁷The inequality $\kappa < \tilde{\kappa}$ is true when k_2 is sufficiently large.

Since, by the choice of k_2 , $|\partial_\zeta G(\zeta; Y; \lambda)| \leq k_0 < 1$, uniformly with respect to $Y \in \mathbb{S}^{n-1} \times \{x \in \mathbb{R}^n : \langle x \rangle \leq \varkappa \lambda\}$, $\lambda \geq \lambda_0$, G has a unique fixed point $\zeta_0^* = \zeta_0^*(Y; \lambda)$, smoothly depending on the parameters, see the Appendix for more details. Since

$$(4.13) \quad \begin{aligned} \partial_t^2 F_2(X, Y; \lambda) &= S_T^{m-1,0}(x, (\lambda \zeta)^{\frac{1}{m}}), \\ \partial_t \partial_\zeta F_2(X, Y; \lambda) &= q_\psi(x, \varsigma) (1 + \zeta_0 \zeta^{-1} (S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}}) + t S_T^{m-1,0}(x, (\lambda \zeta)^{\frac{1}{m}}))), \\ \partial_\zeta^2 F_2(X, Y; \lambda) &= t \zeta^{-1} (S^{-1,1}(x, (\lambda \zeta)^{\frac{1}{m}}) + t S_T^{m-1,1}(x, (\lambda \zeta)^{\frac{1}{m}})), \end{aligned}$$

we can assume that $\lambda \zeta \geq k_2^m$ and the choices of the other parameters imply, on $\text{supp } V_2$,

$$\partial_t^2 F_2(X, Y; \lambda) < \frac{\varepsilon}{2}, \quad \partial_t \partial_\zeta F_2(X, Y; \lambda) \sim \langle x \rangle, \quad \partial_\zeta^2 F_2(X, Y; \lambda) < \frac{\varepsilon}{2} \langle x \rangle^2.$$

So we have proved that, on $\text{supp } V_2$,

$$(4.14) \quad \begin{aligned} M &= F_{2,X}''(X_0^*(Y; \lambda), Y; \lambda) = \begin{pmatrix} M_{11} & M_{12} \\ M_{12} & 0 \end{pmatrix} = \\ &= \begin{pmatrix} S_T^{m-1,0}(x, (\lambda \zeta_0^*)^{\frac{1}{m}}) & q_\psi(x, \varsigma) \left[1 + \frac{\zeta_0}{\zeta_0^*} S^{-1,0}(x, (\lambda \zeta_0^*)^{\frac{1}{m}}) \right] \\ q_\psi(x, \varsigma) \left[1 + \frac{\zeta_0}{\zeta_0^*} S^{-1,0}(x, (\lambda \zeta_0^*)^{\frac{1}{m}}) \right] & 0 \end{pmatrix} \\ &\Rightarrow \det(M) = -q_\psi(x, \varsigma)^2 \left[1 + \frac{\zeta_0}{\zeta_0^*(Y; \lambda)} S^{-1,0}(x, (\lambda \zeta_0^*(Y; \lambda))^{\frac{1}{m}}) \right]^2 \sim \langle x \rangle^2, \quad \|M\| \sim \langle x \rangle. \end{aligned}$$

By (1.1), (4.12), and $\zeta_0^* = G(\zeta_0^*; Y; \lambda)$, $(X, Y) \in S_X \times S_Y \supseteq \text{supp } V_2(\cdot; \lambda)$, we also find

$$|X_0^*(Y; \lambda) - X_0(Y)| = |\zeta_0^*(Y; \lambda) - \zeta_0(Y)| = |\zeta_0 S^{-1,0}(x, (\lambda \zeta_0^*(Y; \lambda))^{\frac{1}{m}})| \leq \frac{A\varepsilon}{2} \langle x \rangle^{-1},$$

uniformly with respect to $\lambda \geq \lambda_0$. The proof is complete. \square

Remark 4.11. The choice of k_2 depends only on the properties of q and on the values of k_1 and ε , that is: we first fix $k_1 > 4AC > 2AC > 2$ and $\varepsilon \in (0, \frac{1}{2})$, then $T > 0$ small enough as explained at the beginning of the proof of Proposition 4.4, then $k_2 > 1$ as explained in the proofs of Propositions 4.8 and 4.10, then, finally, $\lambda_0 > 2k_1 \langle 2k_2 \rangle^m$.

The next Lemma says that the presence in the amplitude of factors which vanish at $X = X_0^*$ implies the gain of negative powers of λ :

Lemma 4.12. Assume $\alpha \in \mathbb{Z}_+^2$, $|\alpha| > 0$,

$$(4.15) \quad W = W(X, Y; \lambda) < V_2(X, Y; \lambda) t^{\alpha_1} \left[W_{\alpha_1+\alpha_2}(X, Y; \lambda) (\zeta - \zeta_0^*(Y; \lambda))^{\alpha_1+\alpha_2} \right]$$

or

$$W = W(X, Y; \lambda) < V_2(X, Y; \lambda) t^{\alpha_1+\alpha_2} \left[W_{\alpha_2}(X, Y; \lambda) (\zeta - \zeta_0^*(Y; \lambda))^{\alpha_2} \right],$$

W is smooth, $W_k(X, Y; \lambda) < \langle x \rangle^k$, $k \in \mathbb{Z}_+$, and has a SG-behaviour as the factors appearing in the expression of V_2 . Then

$$(4.16) \quad \int e^{i\lambda F_2(X, Y; \lambda)} W(X, Y; \lambda) dX = \lambda^{-|\alpha|} \int e^{i\lambda F_2(X, Y; \lambda)} \widetilde{W}(X, Y; \lambda) dX,$$

where \widetilde{W} has the same SG-behaviour, support and x -order⁸ of V_2 .

⁸Including the powers of ζ .

Proof. By arguments similar to those used in the proof of Proposition 4.8, on $\text{supp } W$

$$\partial_{\zeta} F_2(X, Y; \lambda) > \langle x \rangle |t|, \quad \partial_t F_2(X, Y; \lambda) > \langle x \rangle |\zeta - \zeta_0^*(Y; \lambda)|.$$

Assume that the first condition in (4.15) holds. Under the hypotheses, if $\alpha_1 > 0$, we can first insert $e^{i\lambda F_2(X, Y; \lambda)} = L_{\zeta}^{\alpha_1} e^{i\lambda F_2(X, Y; \lambda)}$ in the left hand side of (4.16), where

$$L_{\zeta} = \frac{D_{\zeta}}{\lambda \partial_{\zeta} F_2(X, Y; \lambda)}, \text{ and integrate by parts } \alpha_1 \text{ times. Similarly, if } \alpha_2 > 0, \text{ we}$$

subsequently use $e^{i\lambda F_2(X, Y; \lambda)} = L_t^{\alpha_2} e^{i\lambda F_2(X, Y; \lambda)}$, $L_t = \frac{D_t}{\lambda \partial_t F_2(X, Y; \lambda)}$, and integrate by parts α_2 times. The assertion then follows, remembering that ζ -derivatives of W produce either an additional ζ^{-1} factor or a lowering of the exponent of $\zeta - \zeta_0^*$, and that $\zeta, \zeta_0^* \sim \langle x \rangle^{-1}$ on $\text{supp } W$. The proof in the case that the second condition in (4.15) holds is the same, using first L_{ζ} and then L_t . \square

Proof of Proposition 4.7. Define,

$$\mathcal{Q} = \mathcal{Q}(X, Y; \lambda) = \langle M(X - X_0^*(Y; \lambda), (X - X_0^*(Y; \lambda))),$$

and, for $s \in [0, 1]$,

$$\mathcal{F}_s(X, Y; \lambda) = \mathcal{Q}(X, Y; \lambda) + s\mathcal{G}(X, Y; \lambda),$$

$$\mathcal{G}(X, Y; \lambda) = F_2(X, Y; \lambda) - \mathcal{Q}(X, Y; \lambda).$$

Remembering that $F_2(X_0^*(Y), Y; \lambda) = 0$, $F'_{2,X}(X_0^*(Y), Y; \lambda) = 0$, \mathcal{Q} is the Taylor polynomial of degree two of F_2 at $X = X_0^*$, so that \mathcal{G} vanishes of order 3 at $X = X_0^*$. Obviously, $\mathcal{F}_0(X, Y; \lambda) = \mathcal{Q}(X, Y; \lambda)$ and $\mathcal{F}_1(X, Y; \lambda) = F_2(X, Y; \lambda)$. Write

$$\mathcal{J}_{\tau}(s) = \int e^{i\lambda \mathcal{F}_s(X, Y; \tau^{-1})} V_2(X, Y; \tau^{-1}) dX,$$

$\tau \in (0, \lambda_0^{-1}]$, and consider the Taylor expansion of $\mathcal{J}_{\tau}(s)$ of order $2N - 1$, $N > 1$, so that

$$\left| \mathcal{J}_{\tau}(1) - \sum_{k=0}^{2N-1} \frac{\mathcal{J}_{\tau}^{(k)}(0)}{k!} \right| \leq \sup_{0 < s < 1} \frac{|\mathcal{J}_{\tau}^{(2N)}(s)|}{(2N)!}.$$

Since

$$\mathcal{J}_{\tau}^{(2N)}(s) = (i\lambda)^{2N} \int e^{i\lambda \mathcal{F}_s(X, Y; \tau^{-1})} \mathcal{G}(X, Y; \tau^{-1})^{2N} V_2(X, Y; \tau^{-1}) dX,$$

Remark 4.5 and Lemma 4.12 imply that $|\mathcal{J}_{\tau}^{(2N)}(s)| < \lambda^{-N} \langle x \rangle^{-\frac{N}{m}}$, $\tau \in (0, \lambda_0^{-1}]$, $s \in [0, 1]$: indeed, it is easy to see, by direct computation, that \mathcal{G} can be bounded by linear combinations of expressions of the form

$$t^3, t^2 \left[W_1(X, Y; \tau)(\zeta - \zeta_0^*(Y; \tau)) \right], t \left[W_2(X, Y; \tau)(\zeta - \zeta_0^*(Y; \tau))^2 \right], \\ t \left[W_3(X, Y; \lambda)(\zeta - \zeta_0^*(Y; \tau))^3 \right],$$

with W_k , $k \in \mathbb{Z}_+$, having the required properties. Then, the bound of \mathcal{G}^{2N} will always contain a term of the type $t^{3N} \left[W_{3N}(X, Y; \lambda)(\zeta - \zeta_0^*(Y; \lambda))^{3N} \right]$, which corresponds to the (minimum) value $|\alpha| = 3N$ in (4.15).

Each term $\mathcal{J}_{\tau}^{(k)}(0)$, $k = 0, \dots, 2N - 1$, has the quadratic phase function \mathcal{Q} , which of course also satisfies

$$\partial_{\zeta} \mathcal{Q}(X, Y; \tau^{-1}) > \langle x \rangle |t|, \quad \partial_t \mathcal{Q}(X, Y; \tau^{-1}) > \langle x \rangle |\zeta - \zeta_0^*(Y; \tau^{-1})|.$$

Then, denoting by Γ the Taylor expansion of \mathcal{G} at X_0^* of order $3N$, we observe that $\mathcal{G}^k - \Gamma^k$ can be bounded by polynomial expressions in $X - X_0^*$ of the kind appearing

in the right hand side of (4.15), with $|\alpha| = N + k$ (cfr. the proof of Theorem 7.7.5 in [21]). Setting

$$\mathcal{T}_\tau^k = \int e^{i\lambda Q(X,Y;\tau^{-1})} (i\lambda \Gamma(X,Y;\tau^{-1}))^k V_2(X,Y;\tau^{-1}) dX,$$

Lemma 4.12 implies

$$\mathcal{J}_\tau^{(k)}(0) - \mathcal{T}_\tau^k < \lambda^{-N} \langle x \rangle^{-\frac{m}{m}}.$$

We now apply the Stationary Phase Method to \mathcal{T}_τ^k and prove that

$$(4.17) \quad \mathcal{J}_\tau(1) \sim \sum_{j=0}^{+\infty} d_j(Y;\tau) \lambda^{-1-j},$$

which is a consequence of

$$(4.18) \quad \begin{aligned} \mathcal{T}_\tau^k &\sim \lambda^{-1} \det(M/2\pi i)^{-\frac{1}{2}} \sum_l L_{l,k,Y,\tau} V_2, \\ L_{l,k,Y,\tau} V_2 &= \sum_l (2i\lambda)^{-l} \langle M^{-1} D_X, D_X \rangle^l [(i\lambda \Gamma)^k V_2](X_0^*(Y;\tau^{-1}), Y; \tau^{-1}) / l!, \end{aligned}$$

with M evaluated with τ^{-1} in place of λ . Recalling (4.14), it follows that the inverse matrix M^{-1} satisfies, on $\text{supp } V_2$,

$$M^{-1} = \begin{pmatrix} 0 & \frac{1}{M_{12}} \\ \frac{1}{M_{12}} & -\frac{M_{11}}{M_{12}^2} \end{pmatrix}, \quad \frac{1}{M_{12}} < \langle x \rangle^{-1}, \quad \frac{M_{11}}{M_{12}^2} < \varepsilon \langle x \rangle^{-2}, \quad \|M^{-1}\| \sim \langle x \rangle^{-1},$$

in view of the ellipticity of the involved symbols. Then, the operators $L_{j,k,Y,\tau}$, $j, k \in \mathbb{Z}_+$, do not increase the x -order of the resulting function with respect to that of their arguments, $(i\lambda \Gamma)^k V_2$, which is the same of V_2 , uniformly with respect to τ . The proof of (4.18) then follows by Theorem 7.6.1, the proof of Lemma 7.7.3 and formula (7.6.7) in [21], see also [17, 18]. Indeed, by the mentioned results,

$$\begin{aligned} &\mathcal{J}_\tau^k - \lambda^{-1} \det(M/2\pi i)^{-\frac{1}{2}} \sum_{l \leq k+N} L_{l,k,Y,\tau} V_2 \\ &= \left[\mathcal{J}_\tau^k - \lambda^{-1} \det(M/2\pi i)^{-\frac{1}{2}} \sum_{l \leq k+N+1} L_{l,k,Y,\tau} V_2 \right] + \lambda^{-1} \det(M/2\pi i)^{-\frac{1}{2}} L_{k+N+1,k,Y,\tau} V_2 \\ &< \lambda^{-N-3} \langle x \rangle^{-1} \sum_{|\beta| \leq 2} \|D_X^\beta \langle M^{-1} D_X, D_X \rangle^{k+N+3} [\mathcal{G}^k V_2](X, Y; \tau^{-1})\|_{L^2(\mathbb{R}_X^2)} \\ &\quad + \lambda^{-N-2} \langle x \rangle^{-1} |L_{k+N+1,k,Y,\tau} V_2(X_0^*(Y; \tau^{-1}), Y; \tau^{-1})| \\ &< \lambda^{-N-3} \langle x \rangle^{-1} \left[\int_{c^{-1}\langle x \rangle^{-1}}^{c\langle x \rangle^{-1}} \zeta^{2(\frac{m}{m}-3)} d\zeta \right]^{\frac{1}{2}} + \lambda^{-N-2} \langle x \rangle^{-\frac{m}{m}} \\ &< \lambda^{-N-3} \langle x \rangle^{-\frac{m}{m} + \frac{3}{2}} + \lambda^{-N-2} \langle x \rangle^{-\frac{m}{m}} \\ &< \lambda^{-N-1-\frac{1}{2}} \langle x \rangle^{-\frac{m}{m}}, \quad \lambda \rightarrow +\infty, \end{aligned}$$

since $\langle x \rangle < \lambda$ on $\text{supp } V_2$. It is then enough to sum all the expansions of $\frac{\mathcal{J}_\tau^k}{k!}$, $k = 0, \dots, 2N-1$, and sort the terms by decreasing exponents of λ (as in the proof of Theorem 7.7.5 in [21]) to obtain (4.17) with the usual expression

$$\widetilde{d}_j(Y;\tau) = \det(M/(2\pi i))^{-\frac{1}{2}} \sum_{k-l=j} \sum_{2k \geq 3l} i^{-j} 2^{-k} \langle M^{-1} D_X, D_X \rangle^k [(i\Gamma)^l V_2](X_0^*(Y; \tau^{-1}), Y; \tau^{-1}),$$

so that, in particular,

$$\widetilde{d}_j(Y; \tau) < \langle x \rangle^{-\frac{n}{m}},$$

for any $j \in \mathbb{Z}_+$, $\tau \in (0, \lambda_0^{-1}]$. We can then integrate $\mathcal{J}_\tau(1)$ and its asymptotic expansions with respect to $Y \in \mathbb{S}^{n-1} \times \{x \in \mathbb{R}^n : \langle x \rangle \leq \kappa\lambda\}$ and find

$$J_2(\lambda) = \int_{\langle x \rangle \leq \kappa\lambda} \int_{\mathbb{S}^{n-1}} \mathcal{J}_{\lambda^{-1}}(1) dY \sim \sum_j \lambda^{-1-j} \int_{\langle x \rangle \leq \kappa\lambda} \int_{\mathbb{S}^{n-1}} \widetilde{d}_j(Y; \lambda^{-1}) dY, \quad \lambda \rightarrow +\infty.$$

Let us now focus on the leading coefficient, given by⁹

$$\begin{aligned} \int \widetilde{d}_0(Y; \lambda^{-1}) dY &= \int_{\langle x \rangle \leq \kappa\lambda} \int_{\mathbb{S}^{n-1}} \det(M/(2\pi i))^{-\frac{1}{2}} V_2(X_0^*(Y; \lambda), Y; \lambda) dY \\ &= 2\pi \int_{\langle x \rangle \leq \kappa\lambda} \int_{\mathbb{S}^{n-1}} H_1\left(\frac{\langle x \rangle (\lambda \zeta_0^*(\varsigma, x; \lambda))^{\frac{1}{m}}}{\lambda}\right) H_2((\lambda \zeta_0^*(\varsigma, x; \lambda))^{\frac{1}{m}}) H_3\left(\frac{\zeta_0^*(\varsigma, x; \lambda)}{\zeta_0(\varsigma, x)} - 1\right) \\ &\quad \cdot |\det(M)|^{-\frac{1}{2}} \zeta_0^*(\varsigma, x; \lambda)^{\frac{n}{m}-1} d\varsigma dx \\ &= 2\pi \int_{\langle x \rangle \leq \kappa\lambda} \int_{\mathbb{S}^{n-1}} |\det(M)|^{-\frac{1}{2}} \zeta_0^*(\varsigma, x; \lambda)^{\frac{n}{m}-1} d\varsigma dx, \end{aligned}$$

with M evaluated in $\zeta = \zeta_0^*$. We say that

$$\begin{aligned} \int \widetilde{d}_0(Y; \lambda^{-1}) dY &= 2\pi \int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} \frac{1}{q_\psi(x, \varsigma)^{\frac{n}{m}}} d\varsigma dx + O(\lambda^{\max\{-\frac{1}{m}, n-\frac{n}{m}, -1\}}) \\ &= 2\pi d_0 + O(\lambda^{\max\{-\frac{1}{m}, n-\frac{n}{m}, -1\}}), \quad \lambda \rightarrow +\infty. \end{aligned}$$

To confirm this, first note that $\zeta_0^*(Y; \lambda) \rightarrow \zeta_0(Y)$, $\lambda \rightarrow +\infty$, for any $(Y; \lambda)$ belonging to the support of the integrand, see the Appendix. Moreover, the integrand is uniformly bounded by the summable function $\langle x \rangle^{-\frac{n}{m}}$, and its support is included in the set S . Then, recalling (4.14) and setting $\widetilde{H} = |\zeta_0^2 \det(M)|^{-\frac{1}{2}}$,

$$\begin{aligned} R &= \int_{\langle x \rangle \leq \kappa\lambda} \int_{\mathbb{S}^{n-1}} |\det(M)|^{-\frac{1}{2}} \zeta_0^*(Y; \lambda)^{\frac{n}{m}-1} dY - \int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} \zeta_0(Y)^{\frac{n}{m}-1} dY \\ &= \int_{\langle x \rangle \leq \kappa\lambda} \int_{\mathbb{S}^{n-1}} \zeta_0 \left[\widetilde{H} (\zeta_0^*)^{\frac{n}{m}-1} - \zeta_0^{\frac{n}{m}-1} \right] d\varsigma dx - \int_{\langle x \rangle \geq \kappa\lambda} \int_{\mathbb{S}^{n-1}} \zeta_0^{\frac{n}{m}} d\varsigma dx. \end{aligned}$$

The second integral is always $O(\lambda^{n-\frac{n}{m}})$, since $q_\psi(x, \varsigma) \sim \langle x \rangle$ implies

$$R_2 = \int_{\langle x \rangle \geq \kappa\lambda} \int_{\mathbb{S}^{n-1}} \zeta_0^{\frac{n}{m}} d\varsigma dx \sim \int_{\kappa\lambda}^{+\infty} r^{n-\frac{n}{m}-1} dr = \frac{(\kappa\lambda)^{n-\frac{n}{m}}}{\frac{n}{m} - n}, \quad \lambda \rightarrow +\infty.$$

The first integral can be estimated as follows. Since

$$\zeta_0^* - \zeta_0 = \zeta_0 S^{-1,0}(x, (\lambda \zeta_0^*)^{\frac{1}{m}}) = \zeta_0 O((\lambda \zeta_0^*)^{-\frac{1}{m}}),$$

by the properties of ζ_0^* (see the Appendix) we find

$$\left(\frac{\zeta_0^*}{\zeta_0}\right)^{\frac{n}{m}-1} - 1 = (1 + O((\lambda \zeta_0^*)^{-\frac{1}{m}}))^{\frac{n}{m}-1} - 1 = O((\lambda \zeta_0^*)^{-\frac{1}{m}}) = O(\langle x \rangle^{\frac{1}{m}} \lambda^{-\frac{1}{m}}),$$

⁹Remember that $\psi(0) = 1$ and $a(0, x, \xi) = 1$, for all $x, \xi \in \mathbb{R}^n$. For $\zeta = \zeta_0^*(Y; \lambda)$, the factors H_1, H_2 , and H_3 are identically equal to 1, see the Appendix.

since $S^{-1,0}(x, (\lambda \zeta_0^*)^{\frac{1}{m}}) \ll 1$. By (4.14), we similarly have $\widetilde{H} = 1 + O(\langle x \rangle^{\frac{1}{m}} \lambda^{-\frac{1}{m}})$, so that

$$\begin{aligned} R_1 &= \int_{\langle x \rangle \leq \chi \lambda} \int_{\mathbb{S}^{n-1}} \zeta_0 \left[\widetilde{H}(\zeta_0^*)^{\frac{n}{m}-1} - \zeta_0^{\frac{n}{m}-1} \right] d\zeta dx \\ &= \int_{\langle x \rangle \leq \chi \lambda} \int_{\mathbb{S}^{n-1}} \zeta_0^{\frac{n}{m}} \left[\widetilde{H} \left(\frac{\zeta_0^*}{\zeta_0} \right)^{\frac{n}{m}-1} - 1 \right] d\zeta dx \\ &< \lambda^{-\frac{1}{m}} \int_{\langle x \rangle \leq \chi \lambda} \langle x \rangle^{-\frac{n-1}{m}} dx. \end{aligned}$$

If $n > \frac{1}{1-m} \Leftrightarrow n-1 - \frac{n-1}{m} < -1$, $n \in \mathbb{N}$, $m \in (0, 1)$, the integral in R_1 is convergent for $\lambda \rightarrow +\infty$ and $R_1 = O(\lambda^{-\frac{1}{m}})$. In this case, R_1 contributes an $O(\lambda^{\frac{n}{m}-1-\frac{1}{m}})$ term to the expansion of $I_2(\lambda)$, which is of lower order than the $O(\lambda^{\frac{n}{m}-2})$ term, which is one of the remainders appearing in (4.7). On the other hand, if $n < \frac{1}{1-m}$, the integral in R_1 is divergent, and R_1 itself is $O(\lambda^{n-\frac{n}{m}})$, since, trivially,¹⁰

$$\lim_{\lambda \rightarrow +\infty} \frac{\lambda^{-\frac{1}{m}} \int_0^{\chi \lambda} \frac{r^{n-1}}{(1+r^2)^{\frac{n-1}{2m}}} dr}{\lambda^{n-\frac{n}{m}}} = \lim_{\lambda \rightarrow +\infty} \frac{\int_0^{\chi \lambda} \frac{r^{n-1}}{(1+r^2)^{\frac{n-1}{2m}}} dr}{\lambda^{n-\frac{n-1}{m}}} = \frac{\chi^{n-\frac{n-1}{m}}}{n - \frac{n-1}{m}}.$$

Finally, if $n = \frac{1}{1-m}$, R_1 is $O(\lambda^{-\frac{1}{m}} \ln \lambda)$, by

$$\lim_{\lambda \rightarrow +\infty} \frac{\int_0^{\chi \lambda} \frac{r^{\frac{1}{1-m}-1}}{(1+r^2)^{\frac{1}{2(1-m)}}} dr}{\ln \lambda} = 1,$$

and again contributes a term of lower order than the remainder $O(\lambda^{\frac{n}{m}-2})$. Similar conclusions can be obtained for the subsequent terms of the expansion of $J_2(\lambda)$. The proof is complete, combining the contributions of the remainders like R with the other terms in the expansion of $J_2(\lambda)$, and remembering that

$$I_2(\lambda) = \frac{n}{m} \frac{\lambda^{\frac{n}{m}}}{(2\pi)^n} J_2(\lambda) + O(|\lambda|^{-\infty}) = \frac{n}{m} \lambda^{\frac{n}{m}} \sum_{j=0}^{+\infty} (d_j \lambda^{-1-j} + O(\lambda^{n-\frac{n}{m}-1-j})) + O(|\lambda|^{-\infty}).$$

□

Remark 4.13. The same conclusions concerning the behaviour of R_1 in the final step of the proof of Proposition 4.7 could have been obtained studying the Taylor expansion of the extension of $\zeta_0^*(Y; \tau^{-1})$, $\tau = \lambda^{-1}$, to the interval $[0, \lambda_0^{-1}]_\tau$, similarly to [18].

Proof of Theorem 1.1. The statement for $\mu > m$ follows by the arguments in Section 3 and Propositions 4.2, 4.4, 4.7, summing up the contribution of the local symbol on the exit chart to the contributions of the remaining local symbols, which gives the desired multiple of the integral of $q_\psi^{-\frac{n}{m}}$ on the cosphere bundle as coefficient of the leading term $\lambda^{\frac{n}{m}}$. The remainder has then order equal to the maximum between $\frac{n}{m} - 1$ and n , as claimed. The proof for $\mu < m$ is the same, by exchanging step by step the role of x and ξ . □

¹⁰We observe that the multiplicative constants appearing in the terms $O(\lambda^{n-1})$ in the expansions of $I_1(\lambda)$ and $I_2(\lambda)$ can be estimated by products of expressions in A, C, k_1, ε , times the same power $\chi^{n-\frac{n-1}{m}}$.

APPENDIX

For the sake of completeness, here we illustrate some details of the proof of Theorem 1.1, which we skipped in the previous Sections. They concern, in particular, formula (3.7), which expresses the relation between $\sum_j \widehat{\psi}(\lambda - \eta_j)$ and the oscillatory integrals examined in Section 4. We mainly focus on the aspects which are specific for the manifolds with ends. We also show more precisely how the constants k_1, k_2, λ are involved in the solution of equation (4.12) via the Fixed Point Theorem, completing the proof of Proposition 4.10.

A.1. Solution of Cauchy problems and SG Fourier Integral Operators.

Using the so-called “geometric optics method”, specialised to the pseudodifferential calculus we use (see [7, 8, 9, 10, 11] and [29]), the Cauchy Problem (3.3) can be solved modulo $\mathcal{S}(M)$ by means of an operator family $V(t)$, defined for t in a suitable interval $(-T, T)$, $T > 0$: $V(t)$ induces continuous maps

$$\begin{aligned} V : \mathcal{S}(M) &\rightarrow C^\infty((-T, T), \mathcal{S}(M)), \\ V : \mathcal{S}'(M) &\rightarrow C^\infty((-T, T), \mathcal{S}'(M)) \end{aligned}$$

and

$$(A.1) \quad (D_t - Q) \circ V =: R \in C^\infty((-T, T), L^{-\infty, -\infty}(M)),$$

$$(A.2) \quad Vu|_{t=0} = u, \quad \forall u \in \mathcal{S}'(M).$$

First of all, we recall that the partition of unity $\{\theta_k\}$ and the family of functions $\{\chi_k\}$ of Definition 2.7 can be chosen so that $(\theta_k)_*$ and $(\chi_k)_*$ are SG-symbols of order $(0, 0)$ on U_k , extendable to symbols of the same class defined on \mathbb{R}^n (see [31]).

- Remark A.14.** (1) *The complete symbol of Q depends, in general, on the choice of the admissible atlas, of $\{\theta_k\}$ and of $\{\chi_k\}$. Anyway, if $\{\tilde{q}_k\}$ is another complete symbol of Q , $\kappa(x)(q_k(x, \xi) - \tilde{q}_k(x, \xi)) \in \mathcal{S}(\varphi_k(\Omega_k \cap \tilde{\Omega}_k))$ for an admissible cut-off function κ supported in $\varphi_k(\Omega_k \cap \tilde{\Omega}_k)$.*
- (2) *The solution of (3.3) in the SG-classical case and the properties of φ_k and a_k in (3.5) were investigated in [10] (see also [29], Section 4). In particular, it turns out that $\varphi_k \in C^\infty((-T_k, T_k), S_{cl}^{1,1})$, $T_k > 0$. According to [8], page 101, for every SG phase functions φ of the type involved in the definition of $V(t)$ we also have, for all $x \in \mathbb{R}^n$:*

$$\begin{aligned} |\nabla_\xi \varphi(t; x, \xi) - x| &= |\nabla_\xi \varphi(t; x, \xi) - \nabla_\xi \varphi(0; x, \xi)| = \left| \int_0^t \nabla_\xi \dot{\varphi}(t; x, \xi) dt \right| \\ &= \left| \int_0^t \nabla_\xi (q(x, d_x \varphi(t; x, \xi))) dt \right| \leq C|t|\langle x \rangle, \end{aligned}$$

with a constant $C > 0$ not depending on t, x, ξ . The function $\Phi_{t,\xi}(x) := \nabla_\xi \varphi(t, x, \xi)$ turns out to be a (SG-)diffeomorphism, smoothly depending on the parameters t and ξ (see [7]).

Before proving Theorem 3.4, we state a technical Lemma, whose proof is immediate and henceforth omitted.

Lemma A.15. *Let $U \subset \mathbb{R}^n$ be an open set and define $U_\delta := \bigcup_{x \in U} B(x, \delta\langle x \rangle)$ for arbitrary $\delta > 0$. Assume $\theta, \chi \in C^\infty(\mathbb{R}^n)$ such that $\text{supp } \theta \subset U_{\frac{\delta}{5}}$, $\text{supp } \chi \subset U_\delta$ and $\chi|_{U_{\frac{\delta}{2}}} \equiv 1$. Then, for any diffeomorphism $\Phi_{t,\xi}$, smoothly depending on $t \in (-T, T)$, $\xi \in \mathbb{R}^n$, and such*

that $\forall t, x, \xi$ $|\Phi_{t,\xi}(x) - x| \leq C|t|\langle x \rangle$ with a constant $C > 0$ independent of t, x, ξ ,

$$|t| < \frac{\delta}{4C} \Rightarrow (1 - \chi(x))(\partial^\alpha \theta)(\Phi_{t,\xi}(x)) = 0$$

for any multiindex α and $x, \xi \in \mathbb{R}^n$.

We remark that, since a manifold with ends is, in particular, a SG-manifold, the charts (Ω_k, ψ_k) and the functions $\{\theta_k\}, \{\chi_k\}$, can be chosen such that¹¹

- for a fixed $\delta > 0$, each coordinate open set $U_k = \psi_k(\Omega_k)$, $k = 1, \dots, N$, contains an open subset W_k such that $\bigcup_{x \in W_k} B(x, \delta\langle x \rangle) \subseteq U_k$;
- the supports of θ_k and χ_k , $k = 1, \dots, N$, satisfies hypotheses as the supports of θ and χ in Lemma A.15 (see, e.g., Section 3 of [31] for the construction of functions with the required properties).

Proof of Theorem 3.4. We will write $R \equiv S$ when $R - S \in L^{-\infty, -\infty}(M)$ and $\chi_k \triangleleft \tilde{\chi}_k$ when the functions $\chi_k, \tilde{\chi}_k$ are smooth, non-negative, supported in Ω_k , satisfy $\chi_k \tilde{\chi}_k = \chi_k$ and $(\chi_k)_*, (\tilde{\chi}_k)_*$ are SG-symbols of order $(0, 0)$ on $U_k = \psi_k(\Omega_k)$. Obviously, $R \in L^{-\infty, -\infty}(M)$ implies $R V \in C^\infty((-T, T), L^{-\infty, -\infty}(M))$. To simplify notation, in the computations below we will not distinguish between the functions χ_k, θ_k , etc., and their local representations.

$V(t)$ obviously satisfies (A.2). To prove (A.1), choose functions ζ_k, v_k supported in Ω_k such that $\theta_k \triangleleft \zeta_k \triangleleft \chi_k \triangleleft v_k$. Then $Q \equiv \sum_{k=1}^N \theta_k Q_k \chi_k$ and, for all $k = 1, \dots, N$, $Q \chi_k \equiv v_k Q_k \chi_k$ (see [6], Section 4.4; cfr. also [23]), so that

$$\begin{aligned} Q V(t) &= \sum_{k=1}^N Q \chi_k V_k(t) \theta_k \equiv \sum_{k=1}^N v_k Q_k \chi_k V_k(t) \theta_k \\ &= \sum_{k=1}^N (v_k [Q_k, \chi_k] V_k(t) \theta_k + \chi_k Q_k V_k(t) \theta_k) \\ (A.3) &\equiv \sum_{k=1}^N (v_k [Q_k, \chi_k] \zeta_k V_k(t) \theta_k + v_k [Q_k, \chi_k] (1 - \zeta_k) V_k(t) \theta_k) + D_t V(t) \equiv D_t V(t). \end{aligned}$$

That the first term in the sum (A.3) is smoothing comes from the SG symbolic calculus in \mathbb{R}^n and the observations above, since $\text{sym}([Q_k, \chi_k] \zeta_k) \sim 0$. The same property holds for each k in the second term, provided $t \in I_{T_k}$, $T_k > 0$ small enough. In fact, by Theorems 7 and 8 of [7], $(1 - \zeta_k) V_k(t) \theta_k$ is a SG FIO with the same phase function φ_k and amplitude w_k such that

$$w_k(x, \xi) \sim \sum_{\alpha} \frac{(1 - \zeta_k(x)) (\partial^\alpha \theta_k)(\nabla_\xi \varphi_k(t; \xi, x))}{\alpha!} b_{j\alpha}(t; x, \xi),$$

with suitable SG-symbols $b_{j\alpha}$ defined in terms of φ_k and a_k . By Remark A.14 and Lemma A.15, $w_k \sim 0$ for $|t|$ small enough. The proof that $V(t)$ satisfies (A.1) is completed once we set $T = \min\{T_1, \dots, T_N\}$. The last part of the Theorem can be proved as in [14], Proposition 12.3, since, setting $W(t) := U(-t) V(t)$, it is easy to see $D_t W(t) \equiv 0$, so that $W(0) = I \Rightarrow W(t) \equiv I \Rightarrow V(t) \equiv U(t)$, with smooth dependence on t , as claimed. \square

¹¹Actually, this is relevant only for $k = N$.

A.2. Trace formula and asymptotics for $A \in EL_{cl}^{r,1}(M)$.

Proof of Lemma 3.3. Consider first the finite sum

$$k_J(x, y) = \sum_{j=1}^J \widehat{\psi}(-\eta_k) e_k(x) \overline{e_k(y)}$$

and reduce to the local situation (cfr. Schrohe [31]), via the SG-compatible partition of unity $\{\theta_l\}$ subordinate to the atlas \mathcal{A} , by

$$k_J(x, y) = \sum_{r,s=1}^N \sum_{j=1}^J \widehat{\psi}(-\eta_k) (\theta_r e_k)(x) \overline{(\theta_s e_k)(y)} = \sum_{r,s=1}^N k_R^{rs}(x, y).$$

Then, by $e_k \in \mathcal{S}(M)$ and the fact that $(\theta_r)_* = \theta_r \circ \psi_r^{-1}$ is supported and at most of polynomial growth in U_r , it turns out that we can extend $(\theta_r e_k)_*$ and $(\theta_s e_k)_*$ to elements of $\mathcal{S}(\mathbb{R}^n)$. By an argument similar to the proof of Proposition 1.10.11 in [16] (or by direct estimates of the involved seminorms, as in [14]), $(k_J^{rs})_* \rightarrow (k^{rs})_*$ in $\mathcal{S}(\mathbb{R}^n \times \mathbb{R}^n)$ when $J \rightarrow +\infty$, with $(k^{rs})_*$ kernel of $(\theta_r \widehat{\psi}(-Q) \theta_s)_*$. This proves that $\widehat{\psi}(-Q) = \sum_{r,s=1}^N \theta_r \widehat{\psi}(-Q) \theta_s$ is an operator with kernel $K_\psi(x, y) = \sum_{r,s=1}^N k^{rs}(x, y) \in \mathcal{S}(M \times M)$. \square

The proof of Theorem 3.6 is essentially the one in [14], while the proof of Lemma A.16 comes from [16]: we include both of them here for convenience of the reader.

Proof of Theorem 3.6. Setting $G(\lambda) = \int_{-\infty}^{\lambda} \widehat{\psi}(\tau) d\tau$ and integrating (3.8) in $(-\infty, \lambda)$, we obtain

$$\begin{aligned} \int_{-\infty}^{\lambda} \int \widehat{\psi}(\tau - \eta) dN_Q(\eta) d\tau &= \int \left(\int_{-\infty}^{\lambda} \widehat{\psi}(\tau - \eta) d\tau \right) dN_Q(\eta) \\ (A.4) \quad &= \int G(\lambda - \eta) dN_Q(\eta) = \begin{cases} d_0 \lambda^{\frac{n}{m}} + O(\lambda^{n'}) & \text{for } \lambda \rightarrow +\infty \\ O(|\lambda|^{-\infty}) & \text{for } \lambda \rightarrow -\infty. \end{cases} \end{aligned}$$

Now, observe that

$$\int G(\lambda - \eta) dN_Q(\eta) = \sum_{j=1}^{\infty} G(\lambda - \eta_j) = \sum_{j=1}^{\infty} \int_{-\infty}^{\lambda - \eta_j} \widehat{\psi}(\tau) d\tau = \sum_{j=1}^{\infty} \int H(\lambda - \eta_j - \tau) \widehat{\psi}(\tau) d\tau,$$

where $H(\tau)$ is the Heaviside function. Bringing the series under the integral sign, we can write

$$\begin{aligned} \int G(\lambda - \eta) dN_Q(\eta) &= \int \sum_{j=1}^{\infty} H(\lambda - \eta_j - \tau) \widehat{\psi}(\tau) d\tau = \int N_Q(\lambda - \tau) \widehat{\psi}(\tau) d\tau \\ &= N_Q(\lambda) \int \widehat{\psi}(\tau) d\tau + \int [N_Q(\lambda - \tau) - N_Q(\lambda)] \widehat{\psi}(\tau) d\tau = 2\pi N_Q(\lambda) + R(\lambda), \end{aligned}$$

since $\int \widehat{\psi}(\tau) d\tau = 2\pi\psi(0) = 2\pi$. In view of the monotonicity of N and next Lemma A.16 (cfr. Lemma 4.2.8 of [16]), for $\lambda \geq 1$

$$\begin{aligned} |N_Q(\lambda - \tau) - N_Q(\lambda)| &\leq N_Q(\lambda + |\tau|) - N_Q(\lambda - |\tau|) = \int_{\lambda - |\tau|}^{\lambda + |\tau|} dN_Q(\eta) \\ &= \int_{|\lambda - \eta| \leq |\tau|} dN_Q(\eta) \leq C(1 + |\tau|)^{\frac{n}{m}} (1 + |\lambda|)^{\frac{n}{m} - 1} \leq \tilde{C}(1 + |\tau|)^{\frac{n}{m}} \lambda^{\frac{n}{m} - 1}. \end{aligned}$$

We can then conclude that $R(\lambda) = O(\lambda^{\frac{n}{m}-1})$, $\lambda \geq 1$, since $\widehat{\psi} \in \mathcal{S}$, and this, together with (A.4) and (A.5), completes the proof. \square

Lemma A.16. *Under the hypotheses of Theorem 3.6, there exists a constant $C > 0$ such that for any $K \geq 0$ and any $\lambda \in \mathbb{R}$*

$$\int_{|\lambda-\eta| \leq K} dN_Q(\eta) \leq C(1+K)^{\frac{n}{m}}(1+|\lambda|)^{\frac{n}{m}-1}$$

Proof. Let $h \in (0, \widehat{\psi}(0))$ and $[-K_0, K_0]$ such that $\widehat{\psi}(t) \geq h$ for all $t \in [-K_0, K_0]$. Then, trivially,

$$\int_{|\lambda-\eta| \leq K_0} dN_Q(\eta) \leq h^{-1} \int \widehat{\psi}(\lambda - \eta) dN_Q(\eta).$$

Let us now prove that

$$\int \widehat{\psi}(\lambda - \eta) dN_Q(\eta) \leq C_1(1+|\lambda|)^{\frac{n}{m}-1}.$$

Indeed, this is clear for $\lambda \geq \tilde{C} > 0$ and $\lambda \leq -\tilde{C}$, \tilde{C} suitably large, in view of hypothesis (iii). For $\lambda \in [-\tilde{C}, \tilde{C}]$, choose a constant C_1 so large that $\max_{\lambda \in [-\tilde{C}, \tilde{C}]} \int \widehat{\psi}(\lambda - \eta) dN_Q(\eta) \leq C_1(1 + \tilde{C})^{\frac{n}{m}-1}$. This shows that, for all $\lambda \in \mathbb{R}$,

$$(A.6) \quad \int_{|\lambda-\eta| \leq K_0} dN_Q(\eta) \leq C_2(1+|\lambda|)^{\frac{n}{m}-1}.$$

For arbitrary $K > 0$ there exists $l \in \mathbb{N}$ such that $(l-1)K_0 \leq K < lK_0$. We write

$$\begin{aligned} \int_{|\lambda-\eta| \leq K} dN_Q(\eta) &\leq \int_{|\lambda-\eta| \leq lK_0} dN_Q(\eta) = \sum_{j=0}^{l-1} \int_{jK_0 \leq |\lambda-\eta| \leq (j+1)K_0} dN_Q(\eta) \\ &\leq \sum_{j=0}^{l-1} \left[\int_{\left| \lambda + jK_0 + \frac{K_0}{2} - \eta \right| \leq \frac{K_0}{2}} dN_Q(\eta) + \int_{\left| \lambda - jK_0 - \frac{K_0}{2} - \eta \right| \leq \frac{K_0}{2}} dN_Q(\eta) \right]. \end{aligned}$$

By (A.6), the last sum can be estimated by

$$\begin{aligned} 2C_2 \sum_{j=0}^{l-1} \left(1 + |\lambda| + \left(j + \frac{1}{2} \right) K_0 \right)^{\frac{n}{m}-1} &\leq 2C_2 l \left(1 + |\lambda| + \frac{K_0}{2} + K \right)^{\frac{n}{m}-1} \\ &\leq 2C_2 \left(1 + \frac{K}{K_0} \right) \left(1 + \frac{K_0}{2} + K \right)^{\frac{n}{m}-1} \left(1 + \frac{1}{1 + \frac{K_0}{2} + K} |\lambda| \right)^{\frac{n}{m}-1} \\ &\leq C(1+K)^{\frac{n}{m}}(1+|\lambda|)^{\frac{n}{m}-1}, \end{aligned}$$

as claimed. \square

A.3. The solution $\zeta_0^*(Y; \lambda)$ of the equation $\zeta = G(\zeta; Y; \lambda)$.

We know that $A^{-1}\langle x \rangle^{-1} \leq \zeta_0(\varsigma, x) = q_\psi(x, \varsigma)^{-1} \leq A\langle x \rangle^{-1}$, $Y = (\varsigma, x) \in \widetilde{S}_Y = \mathbb{S}^{n-1} \times \{x \in \mathbb{R}^n : \langle x \rangle \leq \kappa\lambda\}$, and that $k_1 > 4AC > 2AC > 2$. Moreover, $k_2 > 1$ is chosen so large that, in particular, on $\text{supp } U_2 \supset \text{supp } V_2$, the absolute value of the ζ -derivative of G is less than $k_0 \leq 1$, uniformly with respect to $Y \in \widetilde{S}_Y$, $\lambda \geq \lambda_0$, $(X, Y; \lambda) \in \text{supp } V_2$. We want to show that once k_1 is fixed, the choice of such a suitably large $k_2 > 1$ allows to make G a contraction on the compact set $I_x = \left[A^{-1} \left(1 - \frac{\varepsilon}{2} \right) \langle x \rangle^{-1}, A \left(1 + \frac{\varepsilon}{2} \right) \langle x \rangle^{-1} \right] \subset [c^{-1}\langle x \rangle^{-1}, c\langle x \rangle^{-1}]$, uniformly with respect to (ς, x) , $\lambda \geq \lambda_0$, provided $\langle x \rangle \leq \kappa\lambda$,

$\kappa = \left(1 - \frac{\varepsilon}{2}\right)[A(2k_2)^m]^{-1}$. This gives the existence and unicity of $\zeta_0^*(Y; \lambda) \in I_x$ such that $X_0^*(Y; \lambda) = (0, \zeta_0^*(Y; \lambda))$ is the unique stationary point of $F_2(X; Y; \lambda)$, with respect to X , which belongs to the support of $V_2(X; Y; \lambda)$ for $\langle x \rangle \leq \kappa\lambda$.

First of all, the presence of the factors $H_1\left(\frac{\langle x \rangle \langle (\lambda \zeta)^{\frac{1}{m}} \varsigma \rangle^m}{\lambda}\right)$ and $H_2((\lambda \zeta)^{\frac{1}{m}})$ in the expression of U_2 imply $(\lambda \zeta)^{\frac{1}{m}} \geq k_2 \Rightarrow \langle (\lambda \zeta)^{\frac{1}{m}} \rangle \leq (1 + k_2^{-2})^{\frac{1}{2}} (\lambda \zeta)^{\frac{1}{m}}$ and

$$\begin{aligned} (2k_1)^{-1} &\leq \frac{\langle x \rangle \langle (\lambda \zeta)^{\frac{1}{m}} \varsigma \rangle^m}{\lambda} \leq \langle x \rangle (1 + k_2^{-2})^{\frac{m}{2}} \zeta \\ &\Rightarrow [2k_1(1 + k_2^{-2})^{\frac{m}{2}}]^{-1} \leq \langle x \rangle \zeta < \frac{\langle x \rangle \langle (\lambda \zeta)^{\frac{1}{m}} \varsigma \rangle^m}{\lambda} \leq 2k_1 \\ &\Rightarrow \zeta \in [c^{-1} \langle x \rangle^{-1}, c \langle x \rangle^{-1}], \quad c = 2k_1(k_2^{-2} + 1)^{\frac{m}{2}}. \end{aligned}$$

Since $k_1 > 4AC > 2AC$, clearly $I_x \subset [c^{-1} \langle x \rangle^{-1}, c \langle x \rangle^{-1}]$. With an arbitrarily chosen $\varepsilon \in (0, \frac{1}{2})$, take $k_2 > \max\{B, 1\}$ such that $\lambda \zeta > k_2^m$ implies $|S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}})| \leq \frac{\varepsilon}{2}$ and $|\zeta_0 \zeta^{-1} S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}})| \leq k_0 < 1$, which is possible, in view of (1.1) and of the fact that $\zeta_0 \zeta^{-1}$ is bounded on $\text{supp} V_2$. Fix $\lambda \geq \lambda_0 > 2k_1(2k_2)^m$ and $\langle x \rangle \leq \kappa\lambda$. Then, on $\text{supp} V_2$,

$$\begin{aligned} \zeta \in I_x &\Rightarrow \lambda \zeta > \left(1 - \frac{\varepsilon}{2}\right)^{-1} A(2k_2)^m \langle x \rangle A^{-1} \left(1 - \frac{\varepsilon}{2}\right) \langle x \rangle^{-1} = (2k_2)^m > k_2^m \\ &\Rightarrow G(\zeta; Y; \lambda) = \zeta_0(1 + S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}})) \in \left[A^{-1} \left(1 - \frac{\varepsilon}{2}\right) \langle x \rangle^{-1}, A \left(1 + \frac{\varepsilon}{2}\right) \langle x \rangle^{-1}\right] = I_x \\ &\Leftrightarrow G(\cdot; Y; \lambda): I_x \rightarrow I_x. \end{aligned}$$

Since $|\partial_\zeta G(\zeta; Y; \lambda)| = |\zeta_0 \zeta^{-1} S^{-1,0}(x, (\lambda \zeta)^{\frac{1}{m}})| \leq k_0 < 1$, for all $\zeta \in I_x$, $\langle x \rangle \leq \kappa\lambda$, we have proved that for any choice of $Y \in \widetilde{S}_Y$, $\lambda \geq \lambda_0$ as above, $G(\cdot; Y; \lambda)$ has a unique fixed point in $\zeta_0^* = \zeta_0^*(Y; \lambda) \in I_x$, solution of $\zeta = G(\zeta; Y; \lambda)$.

By well-known corollaries of the Fixed Point Theorem for strict contractions on compact subsets of metric spaces, we of course have that ζ_0^* depends smoothly on Y and λ . Moreover, since $\zeta_0^* \in I_x$ for all $Y \in \widetilde{S}_Y$, $\lambda \geq \lambda_0$, obviously $\zeta_0^* \sim \langle x \rangle^{-1}$ and

$$\zeta_0^*(Y; \lambda) = \zeta_0(1 + S^{-1,0}(x, (\lambda \zeta_0^*(Y; \lambda))^{\frac{1}{m}})) \rightarrow \zeta_0(\varsigma, x), \quad \lambda \rightarrow +\infty,$$

pointwise for any (ς, x) . Moreover, by the choices of k_1, k_2 and ε ,

$$\begin{aligned} \frac{\langle x \rangle \langle (\lambda \zeta_0^*(\varsigma, x; \lambda))^{\frac{1}{m}} \rangle^m}{\lambda} &= \left[\frac{\langle x \rangle^{\frac{2}{m}}}{\lambda^{\frac{2}{m}}} + \langle \langle x \rangle \zeta_0^* \rangle^{\frac{2}{m}} \right]^{\frac{m}{2}} > A^{-1} \left(1 - \frac{\varepsilon}{2}\right) > k_1^{-1}, \\ \frac{\langle x \rangle \langle (\lambda \zeta_0^*(\varsigma, x; \lambda))^{\frac{1}{m}} \rangle^m}{\lambda} &< \left[\kappa^{\frac{2}{m}} + \left(A \left(1 + \frac{\varepsilon}{2}\right)\right)^{\frac{2}{m}} \right]^{\frac{m}{2}} = A \left[\left(1 - \frac{\varepsilon}{2}\right)^{\frac{2}{m}} A^{-\frac{4}{m}} (2k_2)^{-2} + \left(1 + \frac{\varepsilon}{2}\right)^{\frac{2}{m}} \right]^{\frac{m}{2}} \\ &< k_1, \\ \langle x \rangle \leq \kappa\lambda &\Leftrightarrow \lambda A^{-1} \left(1 - \frac{\varepsilon}{2}\right) \langle x \rangle^{-1} \geq (2k_2)^m \Rightarrow \lambda \zeta_0^*(\varsigma, x; \lambda) \in [(2k_2)^m, +\infty). \end{aligned}$$

These imply, for any $\varsigma \in \mathbb{S}^{n-1}$, $x \in \mathbb{R}^n$, $\lambda \geq \lambda_0$ such that $\langle x \rangle \leq \kappa\lambda$,

$$H_1\left(\frac{\langle x \rangle \langle (\lambda \zeta_0^*(\varsigma, x; \lambda))^{\frac{1}{m}} \rangle^m}{\lambda}\right) = 1 \quad \text{and} \quad 1 - H_2((\lambda \zeta_0^*(\varsigma, x; \lambda))^{\frac{1}{m}}) = 1.$$

Of course, by the choice of H_3 , for $Y \in \widetilde{S}_Y$, $\lambda \geq \lambda_0$,

$$\zeta_0^* \in I_x \Rightarrow H_3 \left[\frac{\zeta_0^*(\zeta, x; \lambda)}{\zeta_0(\zeta, x)} - 1 \right] = 1.$$

REFERENCES

- [1] U. Battisti and S. Coriasco. Wodzicki Residue for Operators on Manifolds with Cylindrical Ends. *Ann. Global Anal. Geom.*, 40, 2:223–249, 2011.
- [2] P. Boggiatto, E. Buzano, and L. Rodino. *Global Hypoellipticity and Spectral Theory*. Akademie Verlag, Berlin, 1996.
- [3] M. Borsero. Microlocal Analysis and Spectral Theory of Elliptic Operators on Non-compact Manifolds. Tesi di Laurea Magistrale in Matematica, Università di Torino, 2011.
- [4] M. Borsero and S. Coriasco. Eigenvalue Asymptotics of Schrödinger Operators on Manifolds with Ends. In preparation, 2011.
- [5] T. Christiansen and M. Zworski. Spectral Asymptotics for Manifolds with Cylindrical Ends. *Ann. Inst. Fourier, Grenoble*, 45, 1:251–263, 1995.
- [6] H. O. Cordes *The Technique of Pseudodifferential Operators*. Cambridge Univ. Press, 1995.
- [7] S. Coriasco. Fourier Integral Operators in SG classes I. Composition Theorems and Action on SG Sobolev Spaces. *Rend. Sem. Mat. Univ. Pol. Torino*, 57, 4:249–302, 1999.
- [8] S. Coriasco. Fourier Integral Operators in SG classes II. Application to SG Hyperbolic Cauchy Problems. *Ann. Univ. Ferrara*, 47:81–122, 1998.
- [9] S. Coriasco and L. Maniccia. Wave Front Set at Infinity and Hyperbolic Linear Operators with Multiple Characteristics. *Ann. Global Anal. Geom.*, 24:375–400, 2003.
- [10] S. Coriasco and P. Panarese. Fourier Integral Operators Defined by Classical Symbols with Exit Behaviour. *Math. Nachr.*, 242:61–78, 2002.
- [11] S. Coriasco and L. Rodino. Cauchy problem for SG-hyperbolic equations with constant multiplicities. *Ric. di Matematica*, 48, (Suppl.):25–43, 1999.
- [12] J. J. Duistermaat. *Fourier Integral Operators*, volume 130 of *Progress in Mathematics*. Birkhäuser Verlag, Boston, 1996.
- [13] Y. V. Egorov and B.-W. Schulze. *Pseudo-differential Operators, Singularities, Applications*, volume 93 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 1997.
- [14] A. Grigis and J. Sjöstrand. *Microlocal Analysis for Differential Operators*. Cambridge Univ. Press, 1994.
- [15] V. Guillemin. A New Proof of Weyl’s Formula on the Asymptotic Distribution of Eigenvalues. *Adv. in Math.*, 55, 2:131–160, 1985.
- [16] B. Helffer. *Théorie Spectrale pour des Opérateurs Globalement Elliptiques*. Asterisque, 112, 1984.
- [17] B. Helffer and D. Robert. Propriétés Asymptotiques du Spectre d’Opérateurs Pseudodifférentiels sur \mathbb{R}^n . *Comm. Partial Differential Operators*, 7, 7:795–882, 1982.
- [18] B. Helffer and D. Robert. Comportement Asymptotiques Précisé du Spectre d’Opérateurs Globalement Elliptique dans \mathbb{R}^n . *C. R. Acad. Paris, Série I*, 292, 47–50, 1981.
- [19] L. Hörmander. The Spectral Function of an Elliptic Operator. *Acta Math.*, 121:193–218, 1968.
- [20] L. Hörmander. On the Asymptotic Distribution of the Eigenvalues of Pseudodifferential Operators on \mathbb{R}^n . *Ark. Math.*, 17, 2:297–313, 1979.
- [21] L. Hörmander. *The Analysis of Linear Partial Differential Operators I-IV*. Classics in Mathematics. Springer, Berlin, –2009. Reprints of the previous editions –1994.
- [22] V. J. Ivrii. *Microlocal Analysis and Precise Spectral Asymptotics*. Springer-Verlag, Berlin, 1998.
- [23] H. Kumano-go. *Pseudo-Differential Operators*. MIT Press, 1981.
- [24] L. Maniccia and P. Panarese. Eigenvalue Asymptotics for a Class of md-Elliptic ψ do’s on Manifolds with Cylindrical Exits. *Ann. Mat. Pura Appl. (4)*, 181, 3:283–308, 2002.
- [25] L. Maniccia, E. Schrohe, and J. Seiler. Complex Powers of Classical SG-pseudodifferential Operators. *Ann. Univ. Ferrara Sez. VII Sci. Mat.*, 52(2):353–369, 2006.
- [26] R. Melrose. *Geometric scattering theory*. Stanford Lectures. Cambridge University Press, Cambridge, 1995.
- [27] A. Mohammed. Comportement Asymptotique, avec Estimation du Reste, des Valor Propres d’une Classe d’Opérateurs Pseudo-différentiels sur \mathbb{R}^n . *Math. Nachr.*, 14:127–186, 1989.
- [28] F. Nicola. Trace Functionals for a Class of Pseudo-differential Operators in \mathbb{R}^n . *Math. Phys. Anal. Geom.*, 6, 1:89–105, 2003.
- [29] F. Nicola and L. Rodino. SG Pseudo-Differential Operators and Weak Hyperbolicity. *Pliska Stud. Math. Bulgar.*, 15:5–20, 2003.
- [30] C. Parenti. Operatori Pseudodifferenziali in \mathbb{R}^n e Applicazioni. *Ann. Mat. Pura Appl.*, 93:359–389, 1972.

- [31] E. Schrohe. Spaces of Weighted Symbols and Weighted Sobolev Spaces on Manifolds. In *Pseudodifferential operators (Oberwolfach, 1986)*, volume 1256 of *Lecture Notes in Math.*, pages 360–377. Springer, Berlin, 1987.
- [32] B.-W. Schulze. *Boundary Value Problems and Singular Pseudo-differential Operators*. Pure and Applied Mathematics (New York). John Wiley & Sons Ltd., Chichester, 1998.
- [33] M. A. Shubin. *Pseudodifferential Operators and Spectral Theory*. Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1987.
- [34] H. Tamura. Asymptotics Formulas with Sharp Remainder Estimates for Eigenvalues of Elliptic Operators of Second Order. *Duke Math. J.*, 49:87–119, 1982.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO, ITALY

INSTITUT FÜR ANALYSIS, GOTTFRIED WILHELM LEIBNIZ UNIVERSITÄT HANNOVER, GERMANY
E-mail address: `sandro.coriasco@unito.it`

C/O DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO, ITALY